

CropPol: a dynamic, open and global database on crop pollination

Alfonso Allen-Perkins^{1,2}, Ainhoa Magrach^{3,4}, Matteo Dainese⁵, Lucas A. Garibaldi^{6,7}, David Kleijn⁸, Romina Rader⁹, James R. Reilly¹⁰, Rachael Winfree¹⁰, Ola Lundin¹¹, Carley M. McGrady¹², Claire Brittain¹³, David J. Biddinger¹⁴, Derek R. Artz¹⁵, Elizabeth Elle¹⁶, George Hoffman¹⁷, James D. Ellis¹⁸, Jaret Daniels^{18,19}, Jason Gibbs²⁰, Joshua W. Campbell^{18,21}, Julia Brokaw²², Julianna K. Wilson²³, Keith Mason²³, Kimiora L. Ward^{13,24}, Knute B. Gundersen²³, Kyle Bobiwash^{16,20}, Larry Gut²³, Logan M. Rowe²³, Natalie K. Boyle^{15,25}, Neal M. Williams¹³, Neelendra K. Joshi²⁶, Nikki Rothwell²⁷, Robert L. Gillespie²⁸, Rufus Isaacs²³, Shelby J. Fleischer²⁵, Stephen S. Peterson²⁹, Sujaya Rao²², Theresa L. Pitts-Singer¹⁵, Thijs Fijen⁸, Virginie Boreux^{30,31}, Maj Rundlöf³², Blandina Felipe Viana^{33,34}, Alexandra-Maria Klein³¹, Henrik G. Smith^{32,35}, Riccardo Bommarco¹¹, Luísa G. Carvalheiro^{36,37}, Taylor H. Ricketts^{38,39}, Jaboury Ghazoul⁴⁰, Smitha Krishnan^{40,41}, Faye E. Benjamin¹⁰, João Loureiro⁴², Sílvia Castro⁴², Nigel E. Raine⁴³, Gerard Arjen de Groot⁴⁴, Finbarr G. Horgan^{45,46}, Juliana Hipólito⁴⁷, Guy Smagghe⁴⁸, Ivan Meeus⁴⁸, Maxime Eeraerts⁴⁸, Simon G. Potts⁴⁹, Claire Kremen⁵⁰, Daniel García⁵¹, Marcos Miñarro⁵², David W. Crowder⁵³, Gideon Pisanty^{54,55}, Yael Mandelik⁵⁵, Nicolas J. Vereecken⁵⁶, Nicolas Leclercq⁵⁶, Timothy Weekers⁵⁶, Sandra A. M. Lindstrom^{11,32,57}, Dara A. Stanley⁵⁸, Carlos Zaragoza-Trello¹, Charlie C. Nicholson¹³, Jeroen Scheper⁸, Carlos Rad⁵⁹, Evan A.N. Marks⁶⁰, Lucie Mota⁴², Bryan Danforth⁶¹, Mia Park⁶¹, Antônio Diego M. Bezerra⁶², Breno M. Freitas⁶², Rachel E. Mallinger⁶³, Fabiana Oliveira da Silva^{34,64}, Bryony Willcox⁹, Davi L. Ramos⁶⁵, Felipe D. da Silva e Silva⁶⁶, Amparo Lázaro⁶⁷, David Alomar, Miguel A. González-Estévez⁶⁷, Hisatomo Taki⁶⁸, Daniel P. Cariveau⁶⁹, Michael P. D. Garratt⁴⁹, Diego N. Nabaes Jodar⁷, Rebecca I. A. Stewart^{32,36}, Daniel Ariza⁴⁸, Matti Pisman⁴⁸, Elinor M. Lichtenberg^{53,70}, Christof Schüepp⁷¹, Felix Herzog⁷², Martin H. Entling⁷¹, Yoko L. Dupont⁷³, Charles D.

Michener^{74,†}, Gretchen C. Daily⁷⁵, Paul R. Ehrlich⁷⁵, Katherine L.W. Burns⁵⁸, Montserrat Vilà^{1,76}, Andrew Robson⁷⁷, Brad Howlett⁷⁸, Leah Blechschmidt⁴³, Frank Jauker⁷⁹, Franziska Schwarzbach⁷⁹, Maike Nesper⁴⁰, Tim Diekötter⁸⁰, Volkmar Wolters⁷⁹, Helena Castro⁴², Hugo Gaspar⁴², Brian A. Nault⁶¹, Isabelle Badenhauer^{81,82}, Jessica D. Petersen⁸³, Teja Tschardt⁸⁴, Vincent Bretagnolle⁸⁵, D. Susan Willis Chan⁴³, Natacha Chacoff⁸⁶, Georg K.S. Andersson^{32,35}, Shalene Jha⁸⁷, Jonathan F. Colville⁸⁸, Ruan Veldtman⁸⁹, Jeferson Coutinho⁹⁰, Felix J. J. A. Bianchi⁹¹, Louis Sutter⁹², Matthias Albrecht⁷², Philippe Jeanneret⁷², Yi Zou⁹³, Anne L. Averill⁹⁴, Agustin Saez⁹⁵, Amber R. Sciligo⁵⁰, Carlos H. Vergara⁹⁶, Elias H. Bloom⁵³, Elisabeth Oeller⁵³, Ernesto I. Badano⁹⁷, Gregory M. Loeb⁹⁸, Heather Grab⁹⁹, Johan Ekroos³⁵, Vesna Gagic^{11,100}, Saul A. Cunningham¹⁰¹, Jens Åström¹⁰², Pablo Cavigliasso¹⁰³, Alejandro Trillo¹, Alice Classen¹⁰⁴, Alice L. Mauchline⁴⁹, Ana Montero-Castaño⁴³, Andrew Wilby¹⁰⁵, Ben A. Woodcock¹⁰⁶, C. Sheena Sidhu¹⁰⁷, Ingolf Steffan-Dewenter¹⁰⁴, Ioannis N. Vogiatzakis¹⁰⁸, José M. Herrera¹⁰⁹, Mark Otieno¹¹⁰, Mary W. Gikungu¹¹¹, Sarah J. Cusser¹¹², Thomas Nauss¹¹³, Lovisa Nilsson³⁵, Jessica Knapp^{32,114}, Jorge J. Ortega-Marcos¹¹⁵, José A. González¹¹⁵, Juliet L. Osborne¹¹⁴, Rosalind Blanche^{116,†}, Rosalind F. Shaw¹¹⁴, Violeta Hevia¹¹⁵, Jane Stout¹¹⁷, Anthony D. Arthur¹¹⁸, Betina Blochtein^{6,119}, Hajnalka Szentgyorgyi¹²⁰, Jin Li¹²¹, Margaret M. Mayfield¹²², Michał Woyciechowski¹²³, Patrícia Nunes-Silva¹¹⁹, Rosana Halinski de Oliveira¹¹⁹, Steve Henry¹¹⁶, Benno I. Simmons¹²⁴, Bo Dalsgaard¹²⁵, Katrine Hansen¹²⁵, Tuanjit Sritongchuay¹²⁶, Alison D. O'Reilly⁵⁸, Fermín José Chamorro García^{127,128}, Guiomar Nates Parra¹²⁷, Camila Magalhães Pigozo¹²⁹, Ignasi Bartomeus¹

¹ Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Isla de la Cartuja, 41092 Sevilla, Spain

² Departamento de Ingeniería Eléctrica, Electrónica, Automática y Física Aplicada, ETSIDI, Universidad Politécnica de Madrid, 28040 Madrid, Spain.

³ Basque Centre for Climate Change-BC3, Edif. Sede 1, 1º, Parque Científico UPV-EHU, Barrio Sarriena s/n, 48940 Leioa, Spain

⁴ IKERBASQUE, Basque Foundation for Science, María Díaz de Haro 3, 48013 Bilbao, Spain

⁵ Eurac Research, Institute for Alpine Environment

⁶ Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto de Investigaciones en Recursos Naturales, Agroecología y Desarrollo Rural, Río Negro, Argentina

⁷ Universidad Nacional de Río Negro, Instituto de Investigaciones en Recursos Naturales, Agroecología y Desarrollo Rural, Río Negro, Argentina

⁸ Plant Ecology and Nature Conservation Group, Wageningen University & Research, Wageningen, The Netherlands

⁹ School of Environment and Rural Science, University of New England, Armidale, Australia

¹⁰ Department of Ecology, Evolution and Natural Resources, Rutgers University, New Brunswick, NJ 08901, USA

¹¹ Department of Ecology, Swedish University of Agricultural Sciences, SE-750 07 Uppsala, Sweden

¹² Department of Applied Ecology, North Carolina State University, Raleigh, NC 27695, USA

¹³ Department of Entomology and Nematology, University of California Davis, Davis, CA 95616, USA

¹⁴ Department of Entomology, Pennsylvania State University Fruit Research and Extension Center, Biglerville, PA 17307, USA

- ¹⁵ USDA-Agricultural Research Service, Pollinating Insects Research Unit, Logan, UT 84322, USA
- ¹⁶ Department of Biological Sciences, Simon Fraser University, Burnaby, BC, V5A1S6 Canada
- ¹⁷ Department of Crop and Soil Science, Oregon State University, Corvallis, OR 97331, USA
- ¹⁸ Entomology and Nematology Department, University of Florida, Gainesville, FL 32611, USA
- ¹⁹ Florida Museum of Natural History, University of Florida, Gainesville, FL 32611, USA
- ²⁰ Department of Entomology, University of Manitoba, Winnipeg, MB R3T 2N2 Canada
- ²¹ USDA Agricultural Research Service, Northern Plains Agricultural Research Laboratory, Sidney, MT 59270, USA
- ²² Department of Entomology, University of Minnesota, St. Paul, MN 55113, USA
- ²³ Department of Entomology, Michigan State University, East Lansing, MI 48824, USA
- ²⁴ National Park Service, Yosemite National Park, CA 95389, USA
- ²⁵ Department of Entomology, Pennsylvania State University, University Park, PA 16802, USA
- ²⁶ Department of Entomology and Plant Pathology, University of Arkansas, Fayetteville, AR 72701, USA
- ²⁷ Northwest Michigan Horticultural Research Center, Michigan State University, Traverse City, MI 49684, USA
- ²⁸ Agriculture and Natural Resource Program, Wenatchee Valley College, Wenatchee, WA 98801, USA
- ²⁹ AgPollen LLC, 14540 Claribel Road, Waterford, CA 95386, USA
- ³⁰ ETH Zürich - Institute for Terrestrial Ecosystems - Ecosystem Management - Universitaetstrasse 16, 8092 Zurich - Switzerland

- ³¹ University of Freiburg - Chair of Nature Conservation and Landscape Ecology - Tennenbacher Str. 4, 79106 Freiburg, Germany
- ³² Department of Biology, Lund University, SE-223 62 Lund, Sweden
- ³³ Biology Institute, Federal University of Bahia, Salvador, Bahia, Brazil
- ³⁴ National Institute of Science and Technology in Inter and Transdisciplinary Studies in Ecology and Evolution - INCT IN-TREE, Salvador, Bahia, Brazil
- ³⁵ Centre for Environmental and Climate Research, Lund University, S-223 62 Lund, Sweden
- ³⁶ Centre for Ecology, Evolution and Environmental Changes (cE3c), University of Lisbon, Lisbon, Portugal
- ³⁷ Ecology Department, Universidade Federal de Goiás (UFG), Goiânia, Brasil
- ³⁸ Gund Institute for Environment, University of Vermont, Burlington, VT USA 05405
- ³⁹ Rubenstein School for Environment and Natural Resources, University of Vermont, Burlington, VT USA 05405
- ⁴⁰ Department of Environmental Systems Science, ETH Zurich, Universitätsstrasse 16, 8092 Zurich, Switzerland
- ⁴¹ Bioversity International, Bangalore 560 065, India.
- ⁴² FLOWer Lab, Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000-456 Coimbra, Portugal
- ⁴³ School of Environmental Sciences, University of Guelph, Guelph, Ontario, N1G 2W1, Canada
- ⁴⁴ Wageningen Environmental Research, Alterra
- ⁴⁵ EcoLaverna Integral Restoration Ecology, Kildinan, Co. Cork, Ireland
- ⁴⁶ Universidad Católica del Maule, Facultad de Ciencias Agrarias y Forestales, Escuela de Agronomía, Casilla 7-D, Curicó, Chile

⁴⁷ National Institute for Research in the Amazon (INPA), Coordination of Research in Biodiversity – COBIO, 2936 André Araújo Ave, Petrópolis, 69067-375 Manaus, AM, Brazil

⁴⁸ Laboratory of Agrozoology, Department of Plant and Crops, Faculty of Bioscience Engineering, Ghent University, Coupure Links 653, Ghent, Belgium

⁴⁹ Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University of Reading, Reading, RG6 6AR, UK

⁵⁰ Department of Environmental Science, Policy and Management, University of California, Berkeley, 137 Mulford Hall, Berkeley, CA 94720-3114, USA

⁵¹ Universidad de Oviedo y Unidad Mixta de Investigación en Biodiversidad (CSIC-Uo-PA), Spain

⁵² Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Spain

⁵³ Department of Entomology, Washington State University

⁵⁴ Tel Aviv University

⁵⁵ The Hebrew University of Jerusalem

⁵⁶ Agroecology Lab, Université Libre de Bruxelles (ULB), Boulevard du Triomphe CP 264/02, B-1050 Brussels, Belgium.

⁵⁷ Swedish Rural Economy and Agricultural Society, SE-291 09 Kristianstad, Sweden

⁵⁸ School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Ireland

⁵⁹ Composting Research Group UBUCOMP, Universidad de Burgos, Faculty of Sciences, Pl. Misael Bañuelos s/n, 09001 Burgos, Spain

⁶⁰ BETA Technological Center, University of Vic–University of Central Catalonia, Carrer de la Laura 13, 08500 Vic, Catalonia, Spain

⁶¹ Cornell University

⁶² Universidade Federal do Ceará, Centro de Ciências Agrárias, Departamento de Zootecnia, Campus Universitário do Pici, Bloco 808, Caixa Postal 12168, CEP 60356-000 Fortaleza, CE, Brazil

⁶³ University of Florida

⁶⁴ Universidade Federal de Sergipe (UFS)

⁶⁵ University of Brasilia

⁶⁶ Federal Institute of Mato Grosso

⁶⁷ Instituto Mediterráneo de Estudios Avanzados (UIB-CSIC). Global Change Research Group. C/ Miquel Marquès 21, 09190, Esporles, Balearic Islands, Spain.

⁶⁸ Forestry and Forest Products Research Institute, Tsukuba. Ibaraki 305-8687, Japan

⁶⁹ Department of Entomology, University of Minnesota, St. Paul, MN 55108, USA

⁷⁰ Department of Biological Sciences, University of North Texas

⁷¹ iES Landau Institute for Environmental Sciences, University of Koblenz-Landau, Germany

⁷² Agroecology and Environment, Agroscope, Reckenholzstrasse 191, Zurich, CH-8046 Switzerland

⁷³ Dept. of Bioscience, Aarhus University, 8410 Roende, Denmark

⁷⁴ Entomology Division, Natural History Museum, University of Kansas, Lawrence, Kansas 66045, USA

⁷⁵ Center for Conservation Biology, Department of Biology, Stanford University, Stanford, CA 94305, USA

⁷⁶ Department of Plant Biology and Ecology, University of Seville, Sevilla, Spain

- ⁷⁷ Applied Agricultural Remote Sensing Centre (AARSC), University of New England, Armidale, Australia
- ⁷⁸ The New Zealand Institute for Plant and Food Research Ltd
- ⁷⁹ Department of Animal Ecology, Justus Liebig University Giessen, Heinrich-Buff-Ring 26-32, D-35392 Giessen, Germany
- ⁸⁰ Department of Landscape Ecology, Kiel University
- ⁸¹ INRAE, Unité de Recherche Pluridisciplinaire Prairies Plantes Fourragères, F-86600 Lusignan, France
- ⁸² UMR 7372, Centre d'Etudes Biologiques de Chizé, Université de la Rochelle & CNRS, 79360 Villiers en Bois, France
- ⁸³ Minnesota Department of Natural Resources
- ⁸⁴ Agroecology, University of Göttingen, Germany
- ⁸⁵ CEBC-CNRS
- ⁸⁶ Instituto de Ecologia Regional. CONICET UNT
- ⁸⁷ University of Texas at Austin
- ⁸⁸ The Centre for Statistics in Ecology, the Environment and Conservation, Department of Statistical Sciences, University of Cape Town, 7701 Rondebosch, South Africa
- ⁸⁹ South African National Biodiversity Institute
- ⁹⁰ Instituto Federal de Educação, Ciência e Tecnologia da Bahia (IFBA)
- ⁹¹ Farming Systems Ecology, Wageningen University and Research, P.O. Box 430, 6700 AK Wageningen, Netherlands
- ⁹² Plant-Production Systems, Agroscope, Route des Eterpys 18, CH-1964 Conthey, Switzerland

⁹³ Department of Health and Environmental Sciences, Xi'an Jiaotong-Liverpool University
Suzhou, Jiangsu Province P.R.China

⁹⁴ Department of Environmental Conservation, University of Massachusetts, 160 Holdsworth
Way, Amherst, MA 01003-9285, USA

⁹⁵ INIBIOMA (CONICET-Universidad Nacional del Comahue) Bariloche - Rio Negro –
Argentina

⁹⁶ Department of Chemical and Biological Sciences, Universidad de las Américas Puebla,
Cholula, Pue. Mexico

⁹⁷ División de Ciencias Ambientales, Instituto Potosino de Investigación Científica y
Tecnológica, A.C., Mexico

⁹⁸ Department of Entomology, Cornell Agritech, Cornell University

⁹⁹ School of Integrative Plant Science, Cornell University

¹⁰⁰ Queensland Department of Agriculture and Fisheries, Ecosciences Precinct, QLD, 4001,
Australia.

¹⁰¹ Fenner School of Environment and Society, the Australian National University, Canberra,
Australia

¹⁰² Norwegian institute for nature research

¹⁰³ Instituto Nacional de Tecnología Agropecuaria (INTA), Estación Experimental Agropecuaria
Concordia. Programa Nacional Apicultura (PNAPI), Argentina

¹⁰⁴ Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg

¹⁰⁵ Lancaster Environment Centre, Lancaster University, UK

¹⁰⁶ Centre for Ecology and Hydrology, Wallingford, UK

¹⁰⁷ San Mateo Resource Conservation District, California, UK

- ¹⁰⁸ Faculty of Pure and Applied Sciences, Open University of Cyprus, Cyprus
- ¹⁰⁹ Mediterranean Institute for Agriculture, Environment and Development, University of Évora, 7000-651, Évora, Portugal
- ¹¹⁰ Department of Agricultural Resource Management, University of Embu, Kenya
- ¹¹¹ Department of Zoology, National Museums of Kenya, Nairobi, Kenya
- ¹¹² Kellogg Biological Station, Michigan State University
- ¹¹³ Environmental Informatics, Faculty of Geography, University of Marburg
- ¹¹⁴ Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, Cornwall, TR10 9FE, UK
- ¹¹⁵ Social-ecological Systems Laboratory, Department of Ecology, Universidad Autónoma de Madrid, Madrid, Spain
- ¹¹⁶ CSIRO, Australia
- ¹¹⁷ Trinity College Dublin
- ¹¹⁸ Department of Agriculture, Water and the Environment, Australia
- ¹¹⁹ Programa de Pós-Graduação em Ecologia e Evolução da Biodiversidade, Escola de Ciência, Pontifícia Univ Católica do Rio Grande do Sul, Porto Alegre, Brasil
- ¹²⁰ Institute of Botany, Faculty of Biology, Jagiellonian University
- ¹²¹ Data2action, Australia
- ¹²² The University of Queensland, The School of Biological Sciences, Brisbane, Queensland Australia 4072
- ¹²³ Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University
- ¹²⁴ Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Cornwall Campus, Penryn TR10 9FE, UK

¹²⁵ Center for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen, 2100 Copenhagen Ø, Denmark

¹²⁶ Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan Province 666303, China

¹²⁷ Laboratorio de Investigaciones en Abejas (LABUN), Departamento de Biología, Universidad Nacional de Colombia, Sede Bogotá.

¹²⁸ Programa de Pós-graduação em Ecologia e Recursos Naturais, Departamento de Biologia, Universidade Federal do Ceará. Fortaleza-CE, Brazil

¹²⁹ University Jorge Amado, Salvador, Bahia, Brazil

* Correspondence and requests for materials should be addressed to Ignasi Bartomeus (email: nacho.bartomeus@gmail.com).

† Deceased

Seventy five percent of the world's food crops benefit from insect pollination. Hence, there has been increased interest in how global change drivers impact this critical ecosystem service. Because standardized data on crop pollination are rarely available, we are limited in our capacity to understand the variation in pollination benefits to crop yield, as well as to anticipate changes in this service, develop predictions, and inform management actions. Here, we present CropPol, a dynamic, open and global database on crop pollination. It contains measurements recorded from 202 crop studies, covering 3,394 field observations, 2,552 yield measurements (i.e. berry weight, number of fruits and kg per hectare, among others), and 47,752 insect records from 48 commercial crops distributed around the globe. CropPol comprises 32 of the 87 leading global crops and commodities that are pollinator dependent. *Malus domestica* is the most represented crop (32 studies), followed by *Brassica napus* (22 studies), *Vaccinium corymbosum* (13 studies), and *Citrullus lanatus* (12 studies). The most abundant pollinator guilds recorded are honey bees (34.22% counts), bumblebees (19.19%), flies other than Syrphidae and Bombyliidae (13.18%), other wild bees (13.13%), beetles (10.97%), Syrphidae (4.87%), and Bombyliidae (0.05%). Locations comprise 34 countries distributed among Europe (76 studies), Northern America (60), Latin America and the Caribbean (29), Asia (20), Oceania (10), and Africa (7). Sampling spans three decades and is concentrated on 2001-05 (21 studies), 2006-10 (40), 2011-15 (88), and 2016-20 (50). This is the most comprehensive open global data set on measurements of crop flower visitors, crop pollinators and pollination to date, and we encourage researchers to add more datasets to this database in the future. This data set is released for non-commercial use only. Credits should be given to this paper (i.e., proper citation), and the products generated with this database should be shared under the same license terms (CC BY-NC-SA).

Key Words: Agricultural management, bees, crop production, flower visiting insects, pollination, pollinator biodiversity

Introduction

Over 37% of Earth's ice-free land area is directly being used by humans for agriculture or settlements (Klein Goldewijk *et al.*, 2017). In fact, agricultural expansion is the main driver of land use change across the planet (Venter *et al.*, 2016). Along with other human-induced global change drivers, such as global warming and nitrogen deposition, land use change is accelerating extinction rates for most taxonomic groups (MEA, 2005). This biodiversity crisis has led many researchers to investigate how species loss affects nature's contributions to people, the set of benefits humans obtain from nature directly, including crop pollination, water purification, climate regulation, or food production (Díaz *et al.*, 2018).

Crop pollination is a critical contribution of nature to people delivered by multiple species of pollinators, mainly insects (Rader *et al.*, 2016). The annual market value of crop pollination worldwide is estimated to be of US\$235 billion-US\$577 billion (IPBES, 2016), with over 75% of agricultural crops benefiting from animal pollination, mainly insects (Klein *et al.*, 2007), and a global increase in the proportion of land cultivated with pollinator dependent crops (Aizen *et al.*, 2019). Recent meta-analyses have documented the importance of wild bee (Garibaldi *et al.*, 2013) and non-bee pollinators (Rader *et al.*, 2016) for crop production, and the pervasive effects that land-use change has on pollinator populations (Garibaldi *et al.*, 2011; Dainese *et al.*, 2019). However, with 87 pollinator-dependent crops produced worldwide (Klein *et al.*, 2007), we are far from a comprehensive view of how pollination services change across crops and their most important varieties, regions, environmental contexts and through time. For

example, we know that only a fraction of worldwide pollinators are important crop pollination service providers (Kleijn *et al.*, 2015), but the turnover of important pollinators through time and space, even for the same crop, has just started to be explored (Winfree *et al.*, 2018). Similarly, despite clear evidence that crop production can be enhanced by pollinators in both experimental (studies underlying Klein *et al.*, 2007 Appendix 2) and natural (Garibaldi *et al.*, 2013) conditions, pollination levels have rarely been included in predictive models of crop yield (Garibaldi *et al.*, 2020).

One of the main barriers preventing developments in our understanding of global change impacts on nature's contributions to people in general, and on crop pollination in particular, is the lack of standardized datasets that relate the abundance of the providers of nature's contributions, and their final contribution through space and time. In the absence of standardized monitoring programs, compiling comparable datasets collected by different researchers in a decentralized way can allow answering global questions in an efficient way (Bartomeus and Dicks, 2019). Hence, only by compiling the relevant data at the right scales we will be able to advance this field of research by developing predictive models and scenarios for the loss of biodiversity and associated contributions nature affords to people. This is especially relevant as both the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) and the Convention on Biological Diversity (CBD) have called for a better assessment of nature's contributions to people that are directly relevant for policy-making.

Developing predictive models largely hinges on data management practices which facilitate the detection, evaluation and iterative forecasting of changes in ecosystem structure and function (Dietze *et al.*, 2018; White *et al.*, 2018; Yenni *et al.*, 2019). To regularly update models and evaluate forecasts in an open and reproducible fashion, data should be collected frequently

and released as quickly as possible under open licenses (Dietze *et al.*, 2018; White *et al.*, 2018). Furthermore, to support reproducibility and ensure that data can be used easily by a variety of researchers and in multiple modelling approaches, best practices in data structure should be employed for managing and storing collected data (Dietze *et al.*, 2018; White *et al.*, 2018; Yenni *et al.*, 2019). Such practices include the use of open licenses, standard data formats, accompanying metadata, version control, and performing quality control tests, among others (White *et al.*, 2013; Wilson *et al.*, 2014; Hampton *et al.* 2015). Yenni *et al.* (2019) and White *et al.* (2018) provide accessible examples of modern workflows for regularly updated data and near-term iterative forecasting systems, featuring version control (using git and Github), automated data management, and quality control checks (using the testthat R package; Wickham, 2011).

These modern approaches to data management can accelerate ecological research and improve our ability to detect and even predict changes in natural ecosystems instrumental for decision-making, such as their ability to provide nature's contributions to people like crop pollination. Thus, we have compiled CropPol, a dynamic and open database of crop pollination data. The dataset comprises data recorded within 202 different studies on crop pollination: 143 of which were collated through previous meta-analyses (Garibaldi *et al.*, 2015; Kleijn *et al.*, 2015; Garibaldi *et al.*, 2016; Rader *et al.*, 2016; Dainese *et al.*, 2019, Reilly *et al.*, 2020), whereas 34 studies contain unpublished information. Since most of those studies only consider floral visitors contacting the flowers' stigma or anthers during their visit, in this database we use the terms potential pollinators and floral visitors with that meaning (see limitations of this definition in section II.C). We provide data for 3,394 field observations, 2,552 yield measurements, and 47,752 insect records across 48 commercial crops, distributed throughout the globe (see figures

1-5). Furthermore, CropPol comprises 32 of the 87 leading global crops and commodities that benefit from pollination according to Klein *et al.* (2007) (see figure 6). The sampled locations span over 34 countries distributed among Europe (76 studies), Northern America (60), Latin America and the Caribbean (29), Asia (22), Oceania (10), and Africa (7) (figures 1-5). Data collection occurred from 1990 to 2020. CropPol represents a major effort to compile open and standardized measures of the effect of floral visitors on crop production, across different environmental scenarios, and over three decades. However, as with any compilation of data assembled from independent data sources with slightly different objectives and protocols, the researchers using CropPol are encouraged to check carefully which sources are appropriate to answer different questions (see limitations and potential enhancements in section II.C). Nevertheless, despite many factors influencing yield formation, as more data are added to the database in the future, CropPol will help to assess the contribution of managed and wild floral visitors to different crop species, information that is still unclear and is pivotal for managing pollinator ecosystem service.




Figure 1. Distribution of the number of studies and types of crops in CropPol for Americas and the Caribbean. Crop ID's are as follows: *Rubus idaeus* (1), *Fragaria x ananassa* (2), *Coffea arabica* (3), *Coffea canephora* (4), *Prunus dulcis* (5), *Brassica napus* (6), *Vaccinium corymbosum* (7), *Passiflora edulis* (8), *Anacardium occidentale* (9), *Annona muricata* (10), *Annona squamosa* (11), *Bixa orellana* (12), *Gossypium hirsutum* (13), *Malpighia emarginata* (14), *Mangifera indica* (15), *Persea americana* (16), *Macadamia integrifolia* (17), *Prunus avium* (18), *Phaseolus vulgaris* L. (19), *Allium porrum* (20), *Malus domestica* (21), *Pyrus communis* (22), *Vaccinium macrocarpon* (23), *Abelmoschus esculentus* (24), *Cucumis sativus* (25), *Lagenaria siceraria* (26), *Luffa acutangula* (27), *Momordica charantia* (28), *Brassica rapa* (29), *Vaccinium meridionale* (30), *Fagopyrum esculentum* (31), *Citrullus lanatus* (32), *Cucurbita pepo* (33), *Prunus cerasus* (34), *Trifolium pratense* (35), *Helianthus annuus* (36), *Vicia faba* (37), *Psidium guajava* (38), *Actinidia deliciosa* (39), *Cajanus cajan* (40), *Citrus limon* (41), *Citrus paradisi* (42), *Capsicum annuum* (43), *Cucumis melo* (44), *Solanum lycopersicum* (45), *Annona squamosa atemoya* (46), *Coffea arabica/robusta* (47), and *Actinidia chinensis* (48). The dots represent

the centroids of the respective countries (in the case of USA, its dot is located in the geographic center of the contiguous United States).




Figure 2. Distribution of the number of studies and types of crops in CropPol for Europe. Crop ID's are those in figure 1. The dots represent the centroids of the respective countries.




Figure 3. Distribution of the number of studies and types of crops in CropPol for Asia. Crop ID's are those in figure 1. The dots represent the centroids of the respective countries.




Figure 4. Distribution of the number of studies and types of crops in CropPol for Oceania. Crop ID's are those in figure 1. The dots represent the centroids of the respective countries.




Figure 5. Distribution of the number of studies and types of crops in CropPol for Africa. Crop ID's are those in figure 1. The dots represent the centroids of the respective countries.




Figure 6. Number of studies included in CropPol on crops used for human food with an annual production of at least 4,000,000 Metric tonnes (Mt). The production data was collected from the FAO crop production list for the year 2018 (FAOSTAT 2018). The markers represent the impact of pollinators on increasing production according to Klein *et al.* (2007), namely: essential, high, modest, and little (see their characterization in section I.E., Description). In the case of coffee and tropical fruits, the markers summarize the degree of dependence of the following crops: *Coffea arabica* (modest), *Coffea canephora* (high), *Annona spp.* (essential) and *Psidium guajava* (modest).

We aim to maintain and update this database, and researchers are encouraged to add more datasets as explained below.

METADATA

Class I. Data set descriptors

I.A. Data set identity

CropPol, a dynamic and open global database on crop pollination

I.B. Data set identification codes

CropPol_field_level_data.csv

CropPol_sampling_data.csv

CropPol_data_ownership.csv

I.C. Data set description

I.C.1. Principal investigators

Ignasi Bartomeus¹ and Alfonso Allen-Perkins¹.

¹ Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Isla de la Cartuja, 41092 Sevilla, Spain.

I.C.2. Abstract

Seventy five percent of the world's food crops benefit from insect pollination. Hence, there has been increased interest in how global change drivers impact this critical ecosystem service. Because standardized data on crop pollination are rarely available, we are limited in our capacity to understand the variation in pollination benefits to crop yield, as well as to anticipate changes in this service, develop predictions, and inform management actions. Here, we present CropPol, a dynamic, open and global database on crop pollination. It contains measurements recorded from 202 crop studies, covering 3,394 field observations, 2,552 yield measurements (i.e. berry weight, number of fruits and kg per hectare, among others), and 47,752 insect records from 48 commercial crops distributed around the globe. CropPol comprises 32 of the 87 leading

global crops and commodities that are pollinator dependent. *Malus domestica* is the most represented crop (32 studies), followed by *Brassica napus* (22 studies), *Vaccinium corymbosum* (13 studies), and *Citrullus lanatus* (12 studies). The most abundant pollinator guilds recorded are honey bees (34.22% counts), bumblebees (19.19%), flies other than Syrphidae and Bombyliidae (13.18%), other wild bees (13.13%), beetles (10.97%), Syrphidae (4.87%), and Bombyliidae (0.05%). Locations comprise 34 countries distributed among Europe (76 studies), Northern America (60), Latin America and the Caribbean (29), Asia (20), Oceania (10), and Africa (7). Sampling spans three decades and is concentrated on 2001-05 (21 studies), 2006-10 (40), 2011-15 (88), and 2016-20 (50). This is the most comprehensive open global data set on measurements of crop flower visitors, crop pollinators and pollination to date, and we encourage researchers to add more datasets to this database in the future. This data set is released for non-commercial use only. Credits should be given to this paper (i.e., proper citation), and the products generated with this database should be shared under the same license terms (CC BY-NC-SA).

D. Key words

Pollination, crop production, agricultural management, pollinator biodiversity, bees, flower visiting insects

E. Description

CropPol incorporates data from 202 crop pollination studies on 48 commercial crops, collected at 3,394 sites between 1990 and 2020, and distributed throughout the globe (figures 1-5). All the sites represent agricultural landscapes that are highly modified habitats for food production. CropPol includes data on crop yield across 2,552 sites (71.19%), abundance for different species of floral visitors across 2,304 sites (67.88%) and visitation rates to crops by different potential pollinator species across 2,004 sites (59.05%) (see figure 7).




Figure 7. Missing information for the following variables in CropPol_field_level_data.csv: Latitude, longitude, abundance (i.e. number of potential pollinator individuals observed), visitation rate (i.e. number of visits recorded per 100 flowers and hour, unless the variable "visitation_rate_units" in CropPol_field_level_data.csv redefines such units), and yield.

Most of the crops included are pollinator-dependent crops used for human consumption and for which annual production is at least 4×10^6 Metric tonnes (i.e., they are leading global crops and commodities; 73.26% of studies and 65.31% of crops considered) (see figure 6). CropPol also includes raw potential pollinator data for 175 of the studies included (86.63%), which represents 47,752 records of visitors (see CropPol_sampling_data.csv).

In our compilation, according to Klein *et al.* (2007) the impact of potential pollinators on increasing production is *essential* in 26 studies (i.e., production reduction by 90% or more without pollinator activity), *high* in 92 (40 to less than 90% reduction), *modest* in 56 (10 to less

than 40%), *little* in 10 (greater than 0 to less than 10%), and *unknown* (dependence on pollination is known but the contribution of pollinators to crop production is not) in 18. The most represented crop is *Malus domestica* (32 studies), followed by *Brassica napus* (22), *Vaccinium corymbosum* (13), and *Citrullus lanatus* (12).

Overall, 62 studies (30.69%) recorded only bees, whereas 140 studies also targeted additional flower visitors (69.31%). Honey bees were the most abundant pollinator recorded (34.22% of the counts or flower visits in CropPol_sampling_data.csv), followed by bumblebees (19.19%), flies other than Syrphidae and Bombyliidae (13.18%), other wild bees (13.13%), beetles (10.97%), Syrphidae (4.87%), non-bee Hymenoptera (3.07%), Lepidoptera (0.38%), and Bombyliidae (0.05%). Most of the flower visitors recorded have been identified to the species or morphospecies levels (77.71% and 7.58%, respectively). The taxonomic resolution of the remaining visitors is distributed as follows: “family/subfamily/superfamily” (5.69%), “genus/subgenus/tribe” (4.78%), “order/suborder” (4.04%), and “other/unknown” (0.04%). In each global sub-region, the number of sampled records varies greatly. The largest number of flower visitation and count records comes from Western Europe (216,193), followed by Northern Europe (120,754), Southern Europe (98,090), Latin America and the Caribbean (40,973), Northern America (33,904), Eastern Asia (16,649), Australia and New Zealand (16,116), Sub-Saharan Africa (12,875), Southern Asia (10,426), South-eastern Asia (5,370), Eastern Europe (2,320), and Western Asia (656). Although the guild composition of each region varies, bees are the most sampled organisms worldwide, except in Northern Europe (see figure 8): Western Europe (68.1%), Northern Europe (34.4%), Southern Europe (80.3%), Latin America and the Caribbean (89.0%), Northern America (90.9%), Eastern Asia (73.1%), Australia and New Zealand (47.0%), Sub-Saharan Africa (87.9%), Southern Asia (91.3%), South-eastern Asia

(94.7%), Eastern Europe (91.6%), and Western Asia (100%). In Northern Europe the main guild of flower visitors was flies other than Syrphidae and Bombyliidae (54.3%), but this effect is strongly influenced by two studies out of 31 (the percentage of bees and other flies without those studies is 72.7% and 14.5%, respectively).




Figure 8. Proportion of recorded counts in CropPol_sampling_data.csv per guild and geographic area, namely: global region (red) and sub-region (black). The total number of studies by geographic area is shown in brackets.

Finally, in figure 9 we show the spatiotemporal coverage of CropPol. As can be observed, the sampling spans over two decades and concentrates around 2001-05 (21 studies), 2006-10 (40), 2011-15 (88), 2016-20 (50).




Figure 9. Number of studies by year and geographic area, namely: global region (red) and sub-region (black). Circle radii are proportional to the number of studies. The total number of studies by geographic area is shown in brackets.

Class II. Research origin descriptors

II.A. Overall project description

II.A.1 Identity

CropPol, a dynamic and open global database on crop pollination

II.A.2 Originators

Same as in I.C.1. Principal investigators.

II.A.3 Period of Study

Data collection reported in studies occurred from 1990 to 2020. This period of study results from the data collated, after making a general requests for data, and a specific call to the

authors of previous meta-analyses on crop pollination (Garibaldi *et al.*, 2015; Kleijn *et al.*, 2015; Garibaldi *et al.*, 2016; Rader *et al.*, 2016; Dainese *et al.*, 2019, Reilly *et al.*, 2020).

II.A.4 Objectives

Our objectives for compiling these data were to summarize open and standardized measures of (i) crop yield, (ii) abundance for different floral visitor species, and (iii) visitation rates to crops by different groups or species of potential pollinators, across different environmental scenarios; and to identify gaps in geography, crops and varieties.

II.A.5 Abstract

Same as in I.C.2. Abstract.

II.A.6 Source (s) of funding

This research was funded through the 2017-2018 Belmont Forum and BiodivERsA joint call for research proposals, under the BiodivScen ERA-Net COFUND programme, and with the funding organisations AEI, NWO, ECCyT and NSF.

The studies that produced the information compiled in our dataset were funded by grants, scholarships, and fellowships given by several organizations. D.K. was supported by the Dutch Ministry of Economic Affairs (BO-11-011.01-0.51, BO-11-011.01-011). R.R. was supported through the programme Bee Minus to Bee Plus and Beyond: Higher Yields from Smarter, Growth-focused Pollination Systems C11X1309, the Ian Potter Foundation (ref:20160225), a Rural Industries Research and Development Corporation grant for the project “Secure Pollination for More Productive Agriculture (RnD4Profit-15-02-035)” and an Australian Research Council Discovery Early Career Researcher Award DE170101349. H.G.S. was supported by the Swedish research council FORMAS. S.A.M.L. was supported by the Swedish Farmers’ Foundation for Agricultural Research, the Swedish Board of Agriculture. B.F.V. was

supported by MCT/CNPq/CT-AGRO N° 24/2009 Pollinators Research Networks - Process: 556050/2009-6; /CAPES/GEF/FAO/UNEP/FUNBIO; FAPESP/CNPQ/PRONEX N° 020/2009. L.G.C. was supported by the Fundação para Ciência e Tecnologia (FCT) and European Union via the programa operacional regional de Lisboa 2014/2020 (project EUCLIPO-028360) and the Brazilian National Council for Scientific and Technological Development (CNPq. Universal 421668/2018-0; PQ 305157/2018-3). J.G. and S.K. were supported by the Mercator Research Program of the World Food System Centre at ETH Zurich, North-South Centre, ETH Zürich and the Professorship of Ecosystem Management, ETH Zürich. J.L. was supported by the Operational group I9Kiwi – Developing strategies for the sustainability of kiwifruit production through creation of an added value product, funded by PDR2020, the European program INTERREG-SUDOE, project POLL-OLE-GI - Pollinator Protection and Ecosystem Services in SUDOE Region (SOE1/P5/E0129). G.A.d.G. was supported by the Dutch Ministry of Economic Affairs (BO-11-011.01-0.51). F.G.H was funded by The Philippines Department of Agriculture - Bureau of Agricultural Research (DA-BAR). R.B. was supported by the Swedish research council FORMAS. J.H. was supported by Capes and Cnpq. S.G.P. was supported by a grants from EU FP7 (GOCE-CT-2003-506675, ALARM) and BBSRC, Defra, NERC, the Scottish Government and the Wellcome Trust, under the Insect Pollinators Initiative (Sustainable pollination services for UK crops). D.G. was supported by PCIN2014-145-C02-02 (MinECo; EcoFruit project BiodivERsA-FACCE2014-74) and CGL2015-68963-C2-2-R (MinECo/FEDER). M.M. was supported by INIA-RTA2013-00139-C03-01 (MinECo/FEDER). D.C. was supported by USDA NIFA Grant #1003539. Y.M. and his researches were supported in parts by the Israel Ministry of Agriculture Research Grant No. 824-0112-08 and the Israel Science Foundation Research Grant No. 919/09, and the Ministry for Science and Culture of

Lower Saxony Grant No. 11-76-251-99-06/08. J.A. was supported by the Research Council of Norway (225019), Norwegian Environment Agency (2012/16642); C.C.N.: NSF-GRFP. J.S. was supported by 2013–2014 BiodivERsA/FACCEJPI joint call for research proposals (project ECODEAL), European Community’s Seventh Framework Programme (FP7/2007–2013) under Grant Agreement No 244090, STEP Project (Status and Trends of European Pollinators, www.step-project.net). E.M. was supported by European program INTERREG-SUDOE, project POLL-OLE-GI - Pollinator Protection and Ecosystem Services in SUDOE Region (SOE1/P5/E0129). L.M. was supported by Portuguese Foundation for Science and Technology (FCT) - SFRH/BD/116043/2016. B.D. and M.P. were supported by Smith Lever and Hatch Funds administered by Cornell University Agricultural Experiment Station and by a USDA-AFRI grant [USDA 2010-03689, B.N.D., PI]. H.S. was supported by FORMAS grant nr. 2014:00254. R.M. was supported by the Wisconsin Dept of agriculture, trade, and consumer protection. B.K.W. was supported by a PhD scholarship from the University of New England and the Federal Government ‘Rural Research and Development for Profit’ grant for the project “Multi-scale monitoring tools for managing Australian Tree Crops: Industry meets innovation” (RnD4Profit-14-01-008); D.L.R. was supported by the National Council for Scientific and Technological Development (CNPQ). F.D.d.S.S. was supported by the Foundation of Support to Research of Federal District (FAPDF, Brazil - project 9852.56.31658.07042016); M.P.D.G. was supported by a grant from BBSRC, Defra, NERC, the Scottish Government and the Wellcome Trust, under the Insect Pollinators Initiative; G.C.D., P.R.E. and T.H.R. were supported by Summit Foundation. K.L.W.B. was supported by the Irish Research Council-EPA Government of Ireland Postgraduate Scholarship, Eva Crane Trust, National University of Ireland Galway. A.J.R. was supported by a Federal Government ‘Rural Research and Development for Profit’

grant for the project “Multi-scale monitoring tools for managing Australian Tree Crops: Industry meets innovation” (RnD4Profit-14-01-008); B.G.H. was supported through the programme Bee Minus to Bee Plus and Beyond: Higher Yields from Smarter, Growth-focused Pollination Systems C11X1309. F.J. was supported by the Deutsche Bundesstiftung Umwelt (DBU). M.N. was supported by Mercator Research Program of the World Food System Centre at ETH Zurich. H.C. was supported by RENATURE - “Programa Operacional Regional do Centro 2014-2020 (Centro2020) - CENTRO-01-0145-FEDER-000007. H.G. was supported by Operational group I9Kiwi – Developing strategies for the sustainability of kiwifruit production through creation of an added value product, funded by PDR2020. S.C. was supported by CULTIVAR project (CENTRO-01-0145-FEDER-000020), co-financed by Centro 2020, Portugal 2020 and European Union, through ERDF. N.P.C. was supported by CONICET/FUNDACION PROYUNGAS, CONICET/FUNDACION PROYUNGAS, FUNDACION ANTORCHAS; J.F.C. and R.V. were supported by the South African National Biodiversity Institute & GEF. F.O.S. was supported by MCT/CNPq/CT-AGRO N° 24/2009 Pollinators Research Networks - Process: 556050/2009-6; /CAPES/GEF/FAO/UNEP/FUNBIO; FAPESP/CNPQ/PRONEX N° 020/2009. J.G.E.C. was supported by MCT/CNPq/CT-AGRO N° 24/2009 Pollinators Research Networks - Process: 556050/2009-6; /CAPES/GEF/FAO/UNEP/FUNBIO; FAPESP/CNPQ/PRONEX N° 020/2009. L.S., M.A., P.J. were supported by EU FP7. C.H.V. was supported by a grant from Mexico’s Environmental Ministry (SEMARNAT-CONACyT2002-C01-0194) to CV. E.H.B. was supported by USDA NIFA Grant #1003539. J.E. was supported by FORMAS grant nr. 2014:00254. A.T. was supported by the Spanish Ministry of Economy and Competitiveness project FLORMAS (CGL2012-33801) and by the Biodiversa-FACCE project ECODEAL (PCIN-2014-048). A.T. was supported by a Severo-Ochoa predoctoral fellowship (SVP-2013-

067592) and by the Super-B COST Action (FA1307:18100). J.M.H. was supported by the Spanish Ministry of Education and Science through a postdoctoral fellowship ‘Juan de la Cierva’ (FPDI-2013-16335), and by the Portuguese national funding agency for science, research and technology (FCT) (IF/00001/2015). A.C. was financially supported by the German Research Foundation (DFG) within the Research Unit FOR1246. A.M.C. was supported by Food from Thought: Agricultural Systems for a Healthy Planet Initiative (Canada First Research Excellence Fund, grant 000054) and a North American Pollinator Protection Campaign grant 2018. M.O. was supported by a PhD Scholarship from the Felix Trust, UK - 2006 – 2010. N.E.R. was supported by Food from Thought: Agricultural Systems for a Healthy Planet Initiative (Canada First Research Excellence Fund, grant 000054), Ontario Ministry for Agriculture, Food and Rural Affairs (grant 2018-3307), Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant (2015-06783) and as the Rebanks Family Chair in Pollinator Conservation by the Weston Family Foundation. S.J.C. and S.J. were supported by Texas Parks and Wildlife Department, the Army Research Office, and the National Science Foundation. F.J.C.G. and G.N.P. were supported by the Food and Agriculture Organization of the United Nations from the Norwegian Environment Agency for a project on “Building Capacity in the Science-Policy Interface of Pollination Services”. J.K. was supported by the Agriculture and Horticulture Development Board [CP118]. J.J.O.-M. was supported by the European Union FEDER INTERREG SUDOE VB program (Project SOE1/P5/E0129). J.A.G. was supported by the European Union FEDER INTERREG SUDOE VB program (Project SOE1/P5/E0129). J.L.O. and R.F.S. were supported by the Natural Environment Research Council UK [NE/J014680/1]. V.H. was supported by the European Union FEDER INTERREG SUDOE VB program (Project SOE1/P5/E0129). H.S. and M.W. were supported by EU FP7: GOCE-CT-

2003-506675 ALARM. B.I.S. was supported by the Royal Commission for the Exhibition of 1851 Research Fellowship. K.H. was supported by SCIENCE grants: Henrik Tofte Jacobsen's Grant = 15000 DKK; William Demant Fonden = 8500 DKK and Knud Højgaards Fond, 13000 DKK. A.D.O.R. was supported by the Science Foundation Ireland. N.J.V, T.W. and N.L. received financial support from the Walloon Region through a research grant delivered by the Direction générale opérationnelle de l'Agriculture, des Ressources naturelles et de l'Environnement (DGO3) for the "Modèle permaculturel" project on biodiversity in micro-farms, as well as from the FNRS/FWO joint programme "EOS — Excellence Of Science" for the project "CliPS: Climate change and its impact on Pollination Services (project 30947854)". A.S. was supported by the Global Environment Fund, United Nations Environment Program, United Nations Food and Agriculture Organization (GEF/UNEP/FAO) Global Pollination Project, with additional support to the Food and Agriculture Organization of the United Nations from the Norwegian Environment Agency for a project on "Building Capacity in the Science-Policy Interface of Pollination Services", and from the International Fund for Agricultural Development for the development of the sampling protocol. A.-M.K. was funded by the Alexander von Humboldt Foundation with a Feodor Lynen Fellowship and by the German Science foundation (DFG, KL 1849/4-1). Her project was funded by the DFG (Germany Science Foundation) and by the DAAD (German Academic Exchange Programme) to support A.-M.K. C.K. was funded by the Hellmann foundation. B.I.S. was supported by a Royal Commission for the Exhibition of 1851 Research Fellowship. B.M.F - thanks the Project "Conservation and Management of Pollinators for Sustainable Agriculture, through an Ecosystem Approach", which is supported by the Global Environmental Facility Bank (GEF), coordinated by the Food and Agriculture Organization of the United Nations (FAO) with implementation support from the

United Nations Environment Programme (UNEP) and supported in Brazil by the Ministry of Environment (MMA) and Brazilian Biodiversity Fund (Funbio). Also to the National Council for Scientific and Technological Development - CNPq, Brasília-Brazil for financial support to the Brazilian Network of Cashew Pollinators (project # 556042/2009-3) and a Productivity Research Grant (#302934/2010-3). A.D.M.B. thanks a Ph.D scholarship financed by The Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. D.S.W.C and N.E.R were funded by Ontario Ministry of Agriculture, Food and Rural Affairs (OMAFRA) (Grant UofG2015-2466), Ontario Ministry of Environment and Climate Change (MOECC) (Best in Science grant BIS201617-06); Natural Sciences and Engineering Research Council (NSERC) (Discovery Grant 2015-06783); Fresh Vegetable Growers of Ontario (FVGO); and Food from Thought: Agricultural Systems for a Healthy Planet Initiative, by the Canada First Research Excellent Fund (grant 000054). L.B. and N.E.R. were supported and funded by Ontario Ministry of Agriculture, Food and Rural Affairs (OMAFRA) (Grant UofG2015-2466); Ontario Ministry of Environment and Climate Change (MOECC) (Best in Science grant BIS201617-06); Natural Sciences and Engineering Research Council (NSERC) (Discovery Grant 2015-06783); Food from Thought: Agricultural Systems for a Healthy Planet Initiative, by the Canada First Research Excellent Fund (grant 000054); the Weston Family Foundation; Natural Sciences and Engineering Research Council Master's level Canada Graduate Scholarship; Ontario Fruit and Vegetable Conference; Ontario Agricultural College at the University of Guelph; University of Guelph; and the Arrell Food Institute. C.Z.T. was funded by Severo-Ochoa Predoctoral Fellowship (SVP-2014-068580), and IMPLANTIN (CGL201565346R). M.E. was supported by Research Foundation Flanders (FWO) (PhD grant 1S71416N). D.A was funded by Research Foundation Flanders (FWO) (Grant 3G0C4218). Y.L.D. was supported by Aarhus University

Research Foundation. D.N.N.J. was financed by UNRN-CONICET. J.M.H. was funded by funded by the iFCT contract IF/00001/2015 funded by Portuguese National Public Agency for Science, Technology and Innovation (FCT). E.H.B received financial support from NSF Graduate Research Fellowship (Grant 124006-001), USDA Predoctoral Fellowship (Grant 2017-67011-26025), Western SARE Graduate Student Grant (Grant GW15-022). D.W.C and E.H.B. were supported by USDA Organic Transitions Grant (Grant 2014-51106-22096). N.M.W. and K.L.W. were supported by grants from the Coalition for Urban Rural Environmental Stewardship and USDA -National Institute for Food and Agriculture Specialty Crop Research Initiative # 2012-51181-20105.

II.B. Specific subproject description

II.B.1 Site description

CropPol comprises data collected across 12 global subregions, namely: Northern America (60 studies), Western Europe (32), Northern Europe (31), Latin America and the Caribbean (29), Southern Europe (12), Australia and New Zealand (10), South-eastern Asia (9), Sub-Saharan Africa (7), Southern Asia (5), Western Asia (3), Eastern Asia (3), and Eastern Europe (1). We provide latitude and longitude coordinates (in World Geodetic System 1984 datum or WGS 84) for 3,022 out of 3,394 field records (see figure 7). Hence, the context can be extracted for those sites. Locations for other fields were not originally recorded or are protected for privacy reasons. For specific uses they can be obtained upon request to the corresponding data-holder.

Sites are variable, but share the common feature of being highly modified habitats for food production. Management information was provided for 63.7% of the sites, and most of the

crops grew under conventional practices of agricultural intensification (78.7%), followed by organic practices (15.5%), integrated pest management (4.5%) and unmanaged (1.3%). Hence, most of the sites may correspond to monocultures of high-yield varieties, cultivated in medium to large arable fields with medium to high input of mineral fertilizers and pesticides (Tscharntke *et. al*, 2005). Detailed characteristics of the habitats sampled can be accessed for 81.4% of the sites in the corresponding original papers (see variable “Publication” in Table 2, and available DOIs in Table 4).

III.B.2 Experimental or sampling design

All studies measure floral visitor abundances or visitation rates to crop plant species within at least five different crop fields (16.80 ± 21.44). Crop field size ranges from 3×10^{-4} to 84,573 ($549.53 \pm 4,348.36$) hectares with total area sampled within these crop fields ranging from 0.15 to 19,800 m² ($936.85 \pm 2,636.74$ m²). Within each crop field potential pollinators were measured using a variety of techniques (see Research Methods) for a time period ranging from 6 to 2,880 minutes (163.55 ± 186.96 minutes). Flowers sampled per census at each site ranged from 5 to 199,822 flowers ($35,452.84 \pm 162,931.10$ flowers).

In addition, 68.31% of the 202 studies included a measure of crop production or yield, such as kg per hectare or weight per fruit, among others (see variable “yield_units” in Table 2). Furthermore, a subset of such studies also includes measures of yield or production within crop plants subject to different treatments: 19.80% of the studies report results for floral visitor exclusion, whereas 12.87% of them provide values for pollen supplementation.

Detailed characteristics of the sampling design (such as data collection frequency, number of sampling rounds, etc.) are available for 83.16% of the studies in the corresponding original papers (see variable “Publication” in Table 2, and available DOIs in Table 4).

II.B.3 Research methods

CropPol includes 202 studies that assess the effect of flower visitors on crop yield for different crop species collected around the world. The file CropPol_field_level_data.csv includes data on crop yield, floral visitor abundance and visitation rates to crops by different potential pollinator species for 68.32%, 85.15% and 45.54% of the studies, respectively. When available, for each study we mentioned the digital object information (DOI) of the original paper/s (see variable “Publication” in Table 2, and Table 4). Thus, the complete research methodology used in those studies can be accessed. Furthermore, in the case of the studies that provided their sampling raw data (175 studies in CropPol_sampling_data.csv), a brief description of the overall sampling methodology (variable “description”) and the method/s that were used to survey a given site (variable “sampling_method”) were included (92.00% and 98.86%, respectively). Studies predominantly used one sampling method (147 studies), few of them reported 2 methods (26), and 2 studies used three methods. 60 studies collected floral visitor data using “sweep netting”, 58 followed “transect counts”, 53 used “focal observations”, 20 used “pan trap, bee bowl, blue vane trap or pitfall traps”, and 7 used “other” methods.

We provide some metrics already calculated in CropPol by using some general heuristics. Regarding the estimation of richness and abundance in each site, on the one hand, pan-trap data were not taken into account to estimate their values, respectively, if other sampling methods were available. Despite their popularity, pan-traps have a suite of flaws that make them poorly equipped to monitor bees (Portman *et al.*, 2020). On the other hand, the values of richness, abundance and visitation rates for a given site were obtained by aggregating the records of insects observed during the total sampling time. Consequently, in this database richness, abundance and visitation rates do not reflect the mean value of the respective surveys or rounds

in each site, but the total one. When possible, visitation rates were only derived from timed observations to a given number of flowers, and their units were set to [visits per 100 flowers and hour]. Richness data were not calculated in a given study if the percentage of identified species (or morphospecies) was lower than or equal to 75%, or when the data was obtained by using pan-traps. However, other assumptions or metrics can be calculated using CropPol, as the raw data is also available in the database.

To compare the sampling effort among studies and sites, on the one hand, we included two variables in CropPol_field_level_data.csv: “total_samped_area” and “total_sampled_time” (see Table 2). Their values are reported for 63.86% and 55.94% of the 202 studies, respectively. On the other hand, in CropPol_sampling_data.csv the following variables were included to account for sampling effort: “total_samped_area”, “total_sampled_time”, and “total_samped_flowers” (see Table 1). Their values are reported for 64.00%, 69.71%, and 22.29% of the 175 studies, respectively (see their values above, in “II.B.2 Experimental or sampling design”).

Taxonomic resolution for floral visitors was collected from the raw data, when information was available (as is the case of the studies in (Dainese *et al.*, 2019)). Otherwise, we tried to estimate the taxonomic rank of the organisms by using the package taxize in R (Chamberlain *et al.*, 2020) and searching in the Integrated Taxonomic Information System (ITIS) and the NCBI Taxonomy databases. Species taxonomy is provided “as is” by the original data-holders.

The data workflow used to compile CropPol comprised the following stages: 1) Initial data gathering using a common template; 2) data processing; 3) author validation of scripts and data; and 4) final publication (see figure 10). Data gathering stage began in January 2020, after

making a general requests for data, and a specific call to the authors of previous meta-analyses on crop pollination (Garibaldi *et al.*, 2015; Kleijn *et al.*, 2015; Garibaldi *et al.*, 2016; Rader *et al.*, 2016; Dainese *et al.*, 2019, Reilly *et al.*, 2020). The general information on this initiative, data requirements, frequently asked questions, as well as the forms we used to collect the data can be accessed in: <https://www.beeproject.science/croppollination.html>

Raw datasets were processed as soon as we received them. For that reason, data gathering and processing stages overlapped. We transposed raw data to CropPol templates by using R-scripts (R Core Team, 2020) under a version control protocol (i.e. git, <https://git-scm.com/>). During that stage, we fixed transcription and format errors, homogenized information, and prepared automated reports on the transposed datasets (see section III.A.4. Data verification for further detail).

The validation of scripts and data stage began in July 2020 and extended to November 2020. We contacted the corresponding author of each dataset and shared with him/her all the materials collected and produced during the previous stages, along with specific queries. The feedback and corrections we received were used to update and fix the raw materials, R-scripts to process them, and the data in CropPol templates, when needed.

Finally, to compile CropPol we merged those studies that were verified and corrected by the corresponding author, and after performing additional quality checks, published in this data paper. All the process is reproducible and can be tracked at Zenodo ([10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)) We also provide all our code files in the DataS1.zip file.




Figure 10. Data workflow in CropPol. After collecting the raw data, the information is transposed to CropPol templates and checked by using R scripts. The materials gathered during the previous stages are shared with the corresponding authors, along with specific queries. The author's feedback and corrections are used to fix errors. Finally, the verified templates are merged into the main database, and the version number is updated.

II.C. Data Limitations and Potential Enhancements

To properly use CropPol to assess the effects of floral visitors on crop yield, some limitations must be considered. Firstly, it should be noted that, besides successful pollination, many other factors will affect crop yields, such as temperature, water availability and/or crop nutrition. Therefore, depending on the questions addressed by the researchers, CropPol should be combined with other bioclimatic databases.

Secondly, users of CropPol should be aware that the value of this database is that it provides a long term archive for standardized raw data that may otherwise be lost. However, it is beyond the scope of the database in its current form to include all methodological considerations. Those researchers using the data set would be encouraged to check the methodologies of original papers (see Table 4) and to ensure that they meet the criteria of any meta-analyses they may be conducting. These methodological descriptions will help to identify, for instance, how yield was determined. For some crops, CropPol yield data were obtained from experiments conducted on selected plants/trees rather than all plants within fields. Furthermore, in the case of perennial crops, several studies include a single year of field sampling and, hence, interannual variations in resource allocation were not assessed.

Another issue that must be taken into account is that most studies assume that flower visitors observed on crop flowers and touching the reproductive parts of the flower are pollinators that translocate pollen among plants, but determining whether these visitors are effective pollinations requires recording if they actually deposit appropriate pollen loads on flower stigmas. Nevertheless, visitation rate is a good proxy of pollinator function delivered (Vazquez *et al.*, 2005). Indeed, there is a positive correlation between floral visitation frequency and single visit effectiveness in systems where honeybees are absent (Page *et al.*, 2021). To support users, CropPol lists how pollinators and yield were measured and provides several yield measures when available. However, as with any compilation of data assembled from independent data sources with slightly different protocols and objectives, CropPol requires a careful evaluation of which sources are appropriate to answer different questions. For example, sampling effort measures are not available in 44.06% of the studies, and those studies might not be suitable for answering detailed questions. Hence, researchers should filter the appropriate data for their

research goals. We are confident that this database will overcome the putative limitations described above as more data is added over time.

In addition, the majority of data in CropPol is from North America and Western Europe. Therefore, large geographical and crop gaps are found especially in the Southern hemisphere and Africa and Asia in particular. Information on crop varieties is available only on 57.92% of studies (48.38% of sites). Hence, crop variety gaps are also present. This is important because pollinator dependence will vary strongly in horticultural varieties depending on whether the variety is self-compatible or self-incompatible. Nevertheless, since we plan to maintain CropPol as a live dataset where more data will be contributed as it becomes available, we hope to bridge these existing data gaps.

Currently, taxonomy in CropPol_sampling_data.csv (variable “pollinator”) is as provided by the authors. We plan to develop additional tests to curate such data. If any researcher identifies data issues that affect this or other variables, he/she can contact the main investigators by opening GitHub issues and/or via email. The CropPol team will fix the dataset and expand the tested requirements and metadata information, accordingly.

To contribute new datasets, we implemented a modern workflow in CropPol’s GitHub repository (user name: ibartomeus; repository name: OBservData). On the one hand, those users that are familiar with GitHub can follow the workflow A in figure 11, namely: (i) clone the repository; (ii) access the template in the “Template” folder; (iii) fill out the information and save the file in “Your_study_folder” with the name “<author’s name>_“<crop>”_<country>_<year>” (e.g. “John_Doe_Malus_domestica_USA_2020.ods”); (iv) run the R-script “importing_single_file” (if any test fail, a report will be created and the data should be fixed); and (v) pull a request to merge the new data, only once the dataset pass all the automated tests.

On the other hand, for non-GitHub users, we proposed an alternative workflow to contribute new studies (see workflow B in figure 11): (i) access the repository site and download the template in the “Template” folder, (ii) fill out the information and name the file as “<author’s name>_“<crop>”_<country>_<year>”, (iii) open an issue in GitHub to let us know where we can access the filled template; (iv) we will test the template and, if any test fail, we will send an email to the corresponding author, asking him/her to fix his/her data. Once we receive a pull request (workflow A) or data that passes all our tests (workflow B), we will rebuild the database and release a new version of CropPol. Major releases will be deposited permanently at Zenodo (accessible using the same DOI).




Figure 11. Data workflow for collecting new datasets. Workflow A is intended for GitHub users, whereas workflow B is for non-GitHub users. See main text for details on each workflow.

CLASS III. DATA SET STATUS AND ACCESSIBILITY

III.A. Status

III.A.1. Latest update

March 2021

III.A.2. Latest archive date

March 2021

III.A.3. Metadata status

Last update 30 March 2021, version submitted

III.A.4. Data verification

Raw data (collected from different sources) was transposed to CropPol templates by using R-scripts (R Core Team, 2020). During that stage, we corrected any transcription errors and homogenized information. Then we checked the format and values of the different variables by using Testthat (Wickham, 2011). For example, if the data holders provided the latitude and longitude of their orchards/fields/plots, we verified that such locations were in the country that they reported. Then, automated reports on the transposed datasets and their test were prepared with R. In order to check the correctness of the results obtained during the processing stage, we shared with the corresponding authors of each dataset (i) the raw data we received, (ii) the R-scripts (where all the transformations performed on the raw data were recorded), (iii) the resulting files (along with a metadata file that contained the description of the variables), and (iv) the report and some queries. The feedback and corrections we received from the corresponding authors was used to update and fix (i) the raw materials, (ii) R-scripts to process them, and (iii)

the data in CropPol templates, when needed. Finally, to compile CropPol we only merged those studies that were verified and corrected by the corresponding author. All the process is reproducible and can be tracked in our permanent repository (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)). We provide all our code in DataS1.zip file.

III.B. Accessibility

III.B.1 Storage location and medium

The original dataset (v1.1.0) of the CropPol database can be accessed from the ECOLOGY repository. Main upgrades of these datasets will be versioned and deposited in Zenodo (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)).

III.B.2. Contact person

Ignasi Bartomeus¹ (nacho.bartomeus@gmail.com) and Alfonso Allen-Perkins¹ (alfonso.allen.perkins@gmail.com)

¹ Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Isla de la Cartuja, 41092 Sevilla, Spain.

III.B.3. Copyright restrictions

CC BY-NC-SA.

III.B.4. Proprietary restrictions

Please cite this data paper when using the data in bulk, but prioritize citing the original datasets when appropriate (see Table 4).

Citation: Allen-Perkins A., A. Magrach, M. Dainese, L. A. Garibaldi, D. Kleijn, R. Rader, J. R. Reilly, R. Winfree, O. Lundin, C. M. McGrady, C. Brittain, D. J. Biddinger, D. R. Artz, E. Elle, G. Hoffman, J. D. Ellis, J. Daniels, J. Gibbs, J. W. Campbell, J. Brokaw, J. K.

Wilson, K. Mason, K. L. Ward, K. B. Gundersen, K. Bobiwash, L. Gut, L. M. Rowe, N. K. Boyle, N. M. Williams, N. K. Joshi, N. Rothwell, R. L. Gillespie, R. Isaacs, S. J. Fleischer, S. S. Peterson, S. Rao, T. L. Pitts-Singer, T. Fijen, V. Boreux, M. Rundlöf, B. F. Viana, A.-M. Klein, H. G. Smith, R. Bommarco, L. G. Carvalheiro, T. H. Ricketts, J. Ghazoul, S. Krishnan, F. E. Benjamin, J. Loureiro, S. Castro, N. E. Raine, G. A. de Groot, F. G. Horgan, J. Hipólito, G. Smagghe, I. Meeus, M. Eeraerts, S. G. Potts, C. Kremen, D. García, M. Miñarro, D. W. Crowder, G. Pisanty, Y. Mandelik, N. J. Vereecken, N. Leclercq, T. Weekers, S. A. M. Lindstrom, D. A. Stanley, C. Zaragoza-Trello, C. C. Nicholson, J. Scheper, C. Rad, E. A.N. Marks, L. Mota, B. Danforth, M. Park, A. D. M. Bezerra, B. M. Freitas, R. E. Mallinger, F. Oliveira da Silva, B. Willcox, D. L. Ramos, F. D. da Silva e Silva, A. Lázaro, D. Alomar, M. A. González-Estévez, H. Taki, D. P. Cariveau, M. P. D. Garratt, D. N. Nabaes Jodar, R. I. A. Stewart, D. Ariza, M. Pisman, E. M. Lichtenberg, C. Schüepp, F. Herzog, M. H. Entling, Y. L. Dupont, C. D. Michener, G. C. Daily, P. R. Ehrlich, K. L.W. Burns, M. Vilà, A. Robson, B. Howlett, L. Blechschmidt, F. Jauker, F. Schwarzbach, M. Nesper, T. Diekötter, V. Wolters, H. Castro, H. Gaspar, B. A. Nault, I. Badenhauer, J. D. Petersen, T. Tschardt, V. Bretagnolle, D. S. Willis Chan, N. Chacoff, G. K.S. Andersson, S. Jha, J. F. Colville, R. Veldtman, J. Coutinho, F. J. J. A. Bianchi, L. Sutter, M. Albrecht, P. Jeanneret, Y. Zou, A. L. Averill, A. Saez, A. R. Sciligo, C. H. Vergara, E. H. Bloom, E. Oeller, E. I. Badano, G. M. Loeb, H. Grab, J. Ekroos, V. Gagic, S. A. Cunningham, J. Åström, P. Cavigliasso, A. Trillo, A. Classen, A. L. Mauchline, A. Montero-Castaño, A. Wilby, B. A. Woodcock, C. S. Sidhu, I. Steffan-Dewenter, I. N. Vogiatzakis, J. M. Herrera, M. Otieno, M. W. Gikungu, S. J. Cusser, T. Nauss, L. Nilsson, J. Knapp, J. J. Ortega-Marcos, J. A. González, J. L. Osborne, R. Blanche, R. F. Shaw, V. Hevia, J. Stout, A. D. Arthur, B. Blochtein, H. Szentgyorgyi, J. Li, M. M. Mayfield, M.

Woyciechowski, P. Nunes-Silva, R. Halinski de Oliveira, S. Henry, B. I. Simmons, B. Dalsgaard, K. Hansen, T. Sritongchuay, A. D. O'Reilly, F. J. Chamorro García, G. Nates Parra, C. M. Pigozo, I. Bartomeus. CropPol: a dynamic, open and global database on crop pollination. Ecology (volume, issue, year, reference number).

III.B.5. Costs

None.

CLASS IV. DATA STRUCTURAL DESCRIPTORS

IV.A. Data Set File

IV.A.1. Identity

- (1) CropPol_field_level_data.csv
- (2) CropPol_sampling_data.csv
- (3) CropPol_data_ownership.csv

Those data files are provided in the DataS1.zip (see the “Final_Data” subfolder).

IV.A.2. Size

- (1) CropPol_field_level_data.csv: 3,394 sites sampled; 1,854 KB
- (2) CropPol_sampling_data.csv: 47,752 floral visitors records; 16,507 KB
- (3) CropPol_data_ownership.csv: 1,109 records; 247 KB

IV.A.3. Format and storage mode

Data tables formatted as comma-separated values (*.csv)

IV.A.4. Header information

See column descriptions in section IV.B.

IV.A.5. Alphanumeric attributes

Mixed.

IV.A.6. Special characters/fields

Both files CropPol_sampling_data.csv and CropPol_field_level_data.csv contain a column that provides clarifications or comments on the values of other variables (see variable “notes” in Tables 1 and 2).

IV.A.7. Authentication procedures

Same as in III.A.4. Data verification.

IV.B. Variable information

- 1) Site level information
- 2) Insect sampling information
- 3) Data ownership/data holders

IV.C. Data anomalies

If no information is available for a given record, this is indicated as 'NA'. Besides, both files CropPol_sampling_data.csv and CropPol_field_level_data.csv contain a column that provides clarifications or comments on the values of other variables (see variable “notes” in Tables 1 and 2).

CLASS V. SUPPLEMENTAL DESCRIPTORS

V.A. Data acquisition

The current data template that we use for data acquisition can be downloaded from (i) the project site (<https://www.beeproject.science/croppollination.html>), (ii) the CropPoll Zenodo

permanent repository (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)), and (iii) the DataS1.zip (see the “Template” subfolder).

Examples of the completed data forms can be accessed in the the CropPoll Zenodo permanent repository (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)) and in the DataS1.zip file (see the “Datasets Processing” subfolder).

Currently the procedures employed to verify that a data set is error free consist of (i) human review, (ii) automatic data verification as indicated above (III.A.4. Data verification). The datasets collected from now on will be automatically verified as indicated at the end of section II.C. Data Limitations and Potential Enhancements (see the workflow for GitHub and non-GitHub users in Fig. 11).

V.B. Related materials

See Table 4 for a list of publications related with the raw data.

V.C. Computer programs and data-processing algorithms

The algorithms used in deriving, processing, or transforming data can be accessed in the DataS1.zip file and the Zenodo repository (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)).

V.D. Archiving

The data is archived for long-term storage and access in Zenodo (DOI: [10.5281/zenodo.5546600](https://doi.org/10.5281/zenodo.5546600)).

ACKNOWLEDGMENTS

I.B. and A.A.-P. thank Kenna Mackenzie and Sarah S. Greenleaf for their respective contributions to CropPol, and Francisco P. Molina (Seville, Spain) for helping with insect

classification. I.B., L.A.G., D.K., R.W., J.R.R., T.P.M.F., A.A.-P., and A.M. were supported by OBServ Project, funded through the 2017-2018 Belmont Forum and BiodivERsA joint call for research proposals, under the BiodivScen ERA-Net COFUND programme, and with the funding organisations AEI, NWO, ECCyT and NSF.

AUTHOR CONTRIBUTIONS

I.B. and A.A.-P. conceived the idea. A.A.-P. compiled and cleaned the data. M.D., L.A.G., D.K., R.R. and J.R.R. invited the co-authors of their respective meta-analyses to participate in the project and provided the original datasets of those data holders who accepted the invitation. The rest of the authors contributed data. All corresponding authors checked the cleaned data, verified its correctness, and provided corrections, when needed. A.A.-P, I.B. and A.M. wrote the manuscript draft. All the authors discussed and revised earlier versions of the manuscript.

Tables

Table 1. Site level information. Description of the fields related with the site level information – file (1) CropPol_field_level_data.csv

Field	Description	Level or range	Example
study_id	identification code for a given study: Author's name+crop name+country+year	Agustin_Saez_Rubus_idae us_Argentina_2014 ... Yi_Zou_Brassica_napus_ China_2015	Thijs_Fijen_Allium_porru m_Italy_2016

		(n=175)	
site_id	identification code for a site within a study	1 ... zec7 (n=1,802)	Arroyo Claro
pollinator	name of the organism recorded	(Dialictus) sp. D ... Zygoptera_sp. (n=2,887)	Eristalis arbustorum
guild	guild of the pollinator	honeybees bumblebees other_wild_bees syrphids humbleflies other_flies beetles non_bee_hymenoptera lepidoptera other	honeybees
identified_to	taxonomic resolution of the pollinator (whether identification is at the level of species,	class ... Unknow (n=38)	species

	morphospecies, genera, etc).		
sampling_method	method to survey organisms. If multiple methods were used per organism, one independent row is added for each method.	10 censuses of 15 minutes observation to a flowering branch ... transects (n=93)	sweepnet
abundance	number of individuals observed/collected. In the case of performing several censuses (transect walks/plant observations), this field reflects the sum of the individuals collected. When specified in “description”, the values may refer to visitation rates.	0.00000e+00 ... 9808 (n=1,726)	1
total_sampled_area	area sampled during each census at each of the sites (e.g. area	0.15 ... 40700	480

	covered by one transect) in [square meters]. In the cases in which there was more than one sampling area within a site, this variable reflects the sum of their respective areas.	(n=195)	
total_sampled_time	time spent sampling [minutes] each field. In the case in which sites were surveyed multiple times, this variable reflects the sum of their respective durations.	0 ... 161280 (n=165)	60
total_sampled_flowers	number of flowers surveyed at each census (e.g., transect) per site. In the cases in which several censuses were performed, this variable reflects the sum of the respective	5 ... 199822.20 (n=333)	225

	counts.		
description	<p>free text to describe the overall methodology, including the number of temporal replicates per site and what a spatial replicate means in the corresponding study.</p>	<p>10 flowers times 30 min .</p> <p>A group of two to three flowers (rarely one or four) were filmed for 30 min at each site, on three different days during bloom, and resulting in recordings of approx. 225 flower-minutes per site.</p> <p>Exact number of flowers filmed given in field level data file and now used to calculate visitation rates, average under</p> <p>total_sampled_flowers</p> <p>...</p> <p>within one crop field, 3 plots for crop measurements and 12 inventory transects were randomly located. 2 inventory rounds per</p>	<p>3 sampling rounds in one season; one 150m observation transect per plot</p>

		transect (1x morning, 1x afternoon) (n=373)	
notes	free text to add comments on the taxa resolution or any other variables	According to the corresponding author, if there are several pan-trap records for a given species at a given site, it means that such record was identified to a morphospecies level. ... total observation area in square meters, total observation time in minutes (n=61)	includes muscids and drosophila

Table 2. Insect sampling information. Description of the fields related with the insect sampling information – file (2) CropPol_sampling_data.csv

Field	Description	Level or range	Example
study_id	identification code for a given study: Author's	Alejandro_Trillo_Fragari a_ananassa_Spain_2016	Bryony_Willcox_Mangi fera_indica_Australia_2

	name+crop name+country+year	... Yi_Zou_Brassica_napus _China_2015 (n=202)	016
site_id	identification code for a site within a study	1 ... zec7 (n=2,272)	Arroyo Claro
crop	crop latin name	Abelmoschus esculentus ... Vicia faba (n=48)	Helianthus annuus
variety	crop variety name	741 ... Yellow passion fruit (n=193)	Koipesol NAPOLI
management	management system implemented in the field: (1) Organic Agriculture, (2) Integrated pest management, and (3) Other Conventional	organic IPM conventional unmanaged NA	conventional

	Practices (4) unmanaged		
country	country where the crop field is located	Argentina... USA (n=34)	Thailand
latitude	latitude (WGS84) of a given field expressed in degrees [°]	-42.12767 ... 59.86528 (n=1,970)	43.44760
longitude	longitude (WGS84) of a given field expressed in degrees [°]	-123.1979 ... 176.3204 (n=1,959)	8.7155910
X_UTM	Easting planar coordinate of a given field expressed in meters	-4,069,306 ... 4,326,346 (n=368)	677,230
Y_UTM	Northing planar coordinate of a given field expressed in meters	142,490 ... 9,757,262 (n=370)	8,526,182
zone_UTM	the UTM zone number	10	32

	of a given field.	.. SAD 69 24S (n=15)	
sampling_start_m onth	month of the year at the beginning of the sampling period (for example, 1 for January, 2 for February and so on)	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12	2
sampling_end_m onth	month of the year at the end of the sampling period (see description for sampling_start_month)	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12	2
sampling_year	year in which the sampling was carried out	1990 ... 2020 (n=27)	2011-2012
field size	area of the field [hectare]	0.000375 ... 84,573 (n=546)	7.5

yield	yield value of a given field	-1.770894 ... 1,500,000 (n=2,202)	72.548722
yield_units	yield units	average fruit set per 100 flowers ... z-score Seeds produced (n=49)	tonnes per hectare
yield2	secondary yield value	-1.414558 ... 10,386.6 (n=1,477)	213.5790
yield2_units	secondary yield units	%pods produced_pod weight ... z-score Seed set (%) (n=28)	Fruit number on fixed branch length per tree
yield_treatments_ no_pollinators	if the results for yield involve exclosures (e.g., bags, etc.), we fill this column with such results (measured as the first	-2.22144444 ... 1,272.60000000 (n=794)	40.00829587

	unit)		
yield_treatments_ pollen_supplemen t	if the results for yield were obtained by using an additional treatment (e.g., hand-pollination, etc.), we fill this column with such results measured as the first unit)	-1.380536 ... 74,780.40300 (n=657)	30
yield_treatments_ no_pollinators2	if the results for secondary yield involve exclosures (e.g., bags, etc.), we fill this column with such results (second yield unit)	-8.577778 ... 258.62 (n=631)	27.9781746
yield_treatments_ pollen_supplemen t2	if the results for yield were obtained by using an additional treatment (e.g., hand-pollination, etc.), we fill this column with such results. (second yield unit)	-3.38888889 ... 215.29100 (n=546)	87.30599647

fruits_per_plant	average number of fruits per plant [count per plant]	0 ... 12,927.55 (n=199)	774.75685
fruit_weight	average fruit weight [grams per fruit]	0.02930331 ... 8,668.006 (n=710)	1.6675
plant_density	amount of crop plants per unit area of crop field [individuals per square meter]	0.006222222 ... 4,485 (n=156)	2.35
seeds_per_fruit	average number of seeds per fruit [count per fruit]	0 ... 308.5 (n=167)	8.2
seeds_per_plant	average number of seeds per plant or pod [count per plant]	10.5 ... 1,427.24 (n=82)	545.48
seed_weight	average seed weight [grams per 100 seeds]	0.0031 ...	3.985

		81.064 (n=107)	
sampling_richness	method/s to survey organisms that is/are used to estimate richness.	"focal observations" ... "transects + pan trap, bee bowl, blue vane trap, pitfall" (n=11)	"transects + focal observations"
observed_pollinator_richness	number of different pollinator species observed [counts]	0 ... 49 (n=63)	17
other_pollinator_richness	estimated number of different species [counts]	0 ... 164.4062 (n=822)	46.93600
other_richness_estimator_method	method used for estimating "other_pollinator_richness", preferably Chao1.	Chao 1 Chao NA (n=3)	Chao 1
richness_restrictions	free text to describe constraints on	all visitors considered ...	bees and hoverflies

	richness/abundance measurements, such as “only bees”, “only non-managed bees”, etc.	only bees (non-managed bees) (n=15)	
sampling_abundance	method/s to survey organisms that is/are used to estimate abundance.	"focal observations" ... "transects" (n=9)	"sweep net"
abundance	total amount of counts along transect lines [counts]. In the case of performing several transect walks, we indicate the sum of the individuals collected.	0 ... 6,001 (n=544)	1,961
ab_honeybee	total amount of transect counts for honey bees [counts]	0 ... 1,750 (n=397)	237
ab_bombus	total amount of transect counts for bumble bees [counts]	0 ... 1,906	171

		(n=210)	
ab_wildbees	total amount of transect counts for other wild bees [counts]	0 ... 2,697.3 (n=198)	415
ab_syrphids	total amount of transect counts for syrphids [counts]	0 ... 1,782 (n=104)	10
ab_humbleflies	total amount of transect counts for bombyliidae [counts]	0 ... 2 (n=4)	1
ab_other_flies	total amount of transect counts for non syrphid or bombyliida diptera [counts]	0 ... 666 (n=84)	56
ab_beetles	total amount of transect counts for coleoptera [counts]	0 ... 4,861 (n=65)	20

ab_lepidoptera	total amount of transect counts for lepidoptera (butterflies and moths) [counts]	0 ... 452 (n=35)	7
ab_nonbee_hyme noptera	total amount of transect counts for nonbee hymenoptera (sawflies, wasps, ants, etc.) [counts]	0 ... 1,147 (n=59)	59
ab_others	total amount of transect counts that were not included in the previous categories [counts]	0 ... 263 (n=56)	3
total_sampled_area	area sampled during each census at each of the sites (e.g. area covered by one transect) in [square meters]. In the cases in which there was more than one sampling area within a site, this variable reflects the sum of their	0.15 ... 19,800 (n=199)	600

	respective areas.		
total_sampled_time	time spent sampling [minutes] each field. In the case in which sites were surveyed multiple times, this variable reflects the sum of their respective durations.	6 ... 2,880 (n=197)	180
sampling_visitation	method/s to survey organisms that is/are used to estimate visitation rates.	"focal observations" ... "transects" (n=5)	"other"
visitation_rate_units	number of legitimate visits (i.e. contacting reproductive structures) to crop units (flowers, branches, etc.), per unit time. Preferred units: [visits per 100 flowers during one hour].	(average number of) visits per 100 flowers and hour ... visits per unit of time (n=21)	visits per tree and hour

visitation_rate	total visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 10,451.77 (n=1,479)	46.4473684
visit_honeybee	guild (honey bees) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 7,574.678 (n=1,284)	20.11935000
visit_bombus	guild (bumble bees) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 492 (n=584)	4.319706000
visit_wildbees	guild (other wild bees) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 4,251.755 (n=874)	2.374101
visit_syrphids	guild (syrphids) visitation rate to crop units (flowers, branches,etc.) [in the	0 ... 1,980.458 (n=467)	0.394736842

	visitation_rate_units].		
visit_humbleflies	guild (bombyliidae) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 593.7041 (n=26)	0.0007105048
visit_other_flies	guild (non syrphid or bombyliidae diptera) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 607.631 (n=310)	2.0314250839
visit_beetles	guild (coleoptera) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 200 (n=130)	0.7117437722
visit_lepidoptera	guild (lepidoptera: butterflies and moths) visitation rate to crop units (flowers, branches,etc.) [in the	0 ... 229.7873 (n=133)	3.1496062992

	visitation_rate_units].		
visit_nonbee_hymenoptera	guild (nonbee hymenoptera: sawflies, wasps, ants, etc.) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 1,332.724 (n=140)	2.1007727741
visit_others	guild (other) visitation rate to crop units (flowers, branches,etc.) [in the visitation_rate_units].	0 ... 113.5246 (n=108)	0.7812500000
Publication	If published, DOI of the publication (preferred) or article reference, if DOI is not available.	10.1111/1365-2664.12977 ... yield data unpublished (n=88)	10.1098/rspb.2013.2686
Credit	list with all authors who need to be given credit	Agustin Saez/CONICET (Universidad Nacional del Comahue) ...	Christof Schüepp, Felix Herzog and Martin H. Entling

		Yoko L. Dupont, Vibeke Simonsen (n=95)	
Email_contact	email for contacting purposes.	agustinsaez@live.com.ar ... yoko.dupont@bios.au.dk (n=82)	entling@uni-landau.de
notes	comments or clarifications on the values of a given variable	" ab_syrphids would be primarily syrphids, but would also include other flies" ... " yield is pure seed yield (without weeds), yield2 is "normal quality yield" (corrected for seed germination rate) " (n=17)	"total_sampled_area: 800 m2 for honeybees and bumblebees, otherwise 400 m2"

Table 3. Data holders information. Description of the fields related with the data ownership information – file (3) CropPol_data_ownership.csv

Field	Description	Level or range	Example
-------	-------------	----------------	---------

study_id	identification code for a given study: Author's name+crop name+country+year	Alejandro_Trillo_Fragaria_ananassa_Spain_2016 ... Yi_Zou_Brassica_napus_China_2015 (n=202)	Bryony_Willcox_Mangifera_indica_Australia_2016
name	name of the co-author. Co-authors could be people directly involved in collecting the data. The main/corresponding author decides who his/her co-authors are. Please, use one line per co-author.	Agustin Saez ... Yoko L. Dupont (n=185)	Charlie C. Nicholson
affiliation	Co-author affiliation. If a given co-author has several affiliations, please, use one line per affiliation.	Aarhus University, Denmark ... Wageningen Environmental Research, Alterra (n=123)	School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Ireland
email	email address of the co-	[deceased]	freitas@ufc.br

	author	... yoko.dupont@bios.au.dk (n=140)	
role	One of the following role categories: (1) Lead author/Corresponding author, (2) Co- author/Co-owner	Lead author/Corresponding author Co-author/Co-owner	Co-author/Co-owner
funding	Funding sources (grants, scholarships, projects, etc.) that supported the co-author	"2013 2014 BiodivERsA FACCEJPI joint call for research proposals (project ECODEAL)" ... "Wisconsin Dept of agriculture, trade, and consumer protection" (n=71)	This study was financially supported by the German Research Foundation (DFG) within the Research Unit FOR1246

Table 4. List of publications related with the raw data.

Publication (DOI)	Study identifier (study_id)
10.1126/science.aac7287	Agustin_Saez_Rubus_idaeus_Argentina_2014, Breno_M_Freitas_Anacardium_occidentale_Brazil_2011,

	<p>Guioamar_Nates_Parra_Vaccinium_meridionale_Colombia_2013,</p> <p>Jens_Astrom_Malus_domestica_Norway_2013,</p> <p>Jens_Astrom_Trifolium_pratense_Norway_2013,</p> <p>Jens_Astrom_Trifolium_pratense_Norway_2014,</p> <p>Ruan_Veldtman_Helianthus_annuus_South_Africa_2011</p>
10.1016/j.baae.2018.05.008	Alejandro_Trillo_Fragaria_ananassa_Spain_2016
10.1098/rspb.2002.2306	Alexandra_Maria_Klein_Coffea_arabica_Indonesia_2000_2001
10.1046/j.1365-2664.2003.00847.x	Alexandra_Maria_Klein_Coffea_canephora_Indonesia_2000_2001
10.1111/j.1365-2664.2012.02144.x	Alexandra_Maria_Klein_Prunus_dulcis_USA_2008
10.1038/ncomms8414	<p>Alexandra_Maria_Klein_Prunus_dulcis_USA_2009,</p> <p>David_Kleijn_Allium_porrum_Italy_2012,</p> <p>Mia_Park_Malus_domestica_USA_2009,</p> <p>Mia_Park_Malus_domestica_USA_2010,</p> <p>Mia_Park_Malus_domestica_USA_2011,</p> <p>Rachael_Winfree_Malus_Domestica_USA_2004,</p> <p>Ruan_Veldtman_Malus_domestica_South_Africa_2011</p>
<p>10.1098/rspb.2013.3148,</p> <p>10.5281/zenodo.12540</p>	Alice_Classen_Coffea_arabica_Tanzania_2011_2012
<p>10.1016/j.agee.2018.05.004,</p> <p>10.1016/j.agee.2019.02.009</p>	<p>Amparo_Lazaro_Prunus_dulcis_Spain_2015,</p> <p>Amparo_Lazaro_Prunus_dulcis_Spain_2016</p>
10.1590/1519-6984.02213	Betina_Blochtein_Brassica_napus_Brazil_2011

10.1111/j.1461-0248.2011.01669.x	Blande_Viana_Passiflora_edulis_Brazil_2005
10.1126/science.1230200	Breno_M_Freitas_Anacardium_occidentale_Brazil_2012, Breno_M_Freitas_Gossypium_hirsutum_Brazil_2011
10.1073/pnas.1517092112	Breno_M_Freitas_Annona_squamosa_Brazil_2013, Breno_M_Freitas_Malpighia_emarginata_Brazil_2011
10.1126/sciadv.aax0121	Breno_M_Freitas_Bixa_orellana_Brazil_2007
10.1038/s41598-019-49535-w	Bryony_Willcox_Mangifera_indica_Australia_2016
10.1038/s41598-019-49535-w, yield data unpublished	Bryony_Willcox_Persea_americana_Australia_2015, Bryony_Willcox_Persea_americana_Australia_2016, Bryony_Willcox_Macadamia_integrifolia_Australia_2016, Bryony_Willcox_Mangifera_indica_Australia_2016_2, Bryony_Willcox_Persea_americana_Australia_2017
10.1016/j.agee.2008.08.001	Carlos_H_Vergara_Coffea_arabica_Mexico_2004
10.1016/j.agee.2018.10.018, 10.1016/j.agee.2017.08.030	Charlie_Nicholson_Vaccinium_corymbosum_USA_2014, Charlie_Nicholson_Vaccinium_corymbosum_USA_2015, Charlie_Nicholson_Vaccinium_corymbosum_USA_2013
10.1098/rspb.2013.2667	Christof_Schuepps_Prunus_avium_Switzerland_2011
10.1111/1365-2664.12060	Dara_St Stanley_Brassica_napus_Ireland_2009
10.1007/s10841-013-9599-z, 10.1007/s11258-014-0301-7	Dara_St Stanley_Brassica_napus_Ireland_2010

10.1371/journal.pone.0204460	Davi_L_Ramos_Phaseolus_vulgaris L_Brazil_2015_2016
10.1093/aesa/88.3.334	David_Kleijn_Vaccinium_macrocarpon_USA_1990, David_Kleijn_Vaccinium_macrocarpon_USA_1991
10.1371/journal.pone.0025172	Dupont_redclover_Denmark_2008, Dupont_redclover_Denmark_2009
10.1016/j.agee.2017.01.031	Eeraerts_etal_sweetcherry_Belgium_2015
10.1016/j.agee.2019.106586	Eeraerts_etal_sweetcherry_Belgium_2016, Eeraerts_etal_sweetcherry_Belgium_2017
10.1126/science.aac7287, 10.26786/1920-7603%282014%2926	Fabiana_Oliveira_da_Silva_Malus_domestica_Brazil_2010, Fabiana_Oliveira_da_Silva_Malus_domestica_Brazil_2011, Fabiana_Oliveira_da_Silva_Malus_domestica_Brazil_2012
10.1007/s10980-009-9331-2	Frank_Jauker_Brassica_napus_Germany_2006
10.1371/journal.pone.0031599	Georg_Andersson_Fragaria_ananassa_Sweden_2009
10.1016/j.agee.2009.05.001	Hajnalka_Szentgyorgyi_Fagopyrum_esculentum_Poland_2005, Simon_Potts_Vicia_faba_UK_2005
10.1016/j.agee.2015.05.004	Heather_Lee_Grab_Fragaria_ananassa_USA_2012
10.1111/j.1744-7348.2009.00326.x, 10.1016/j.baae.2010.08.004	Hisatomo_Taki_Fagopyrum_esculentum_Japan_2007, Hisatomo_Taki_Fagopyrum_esculentum_Japan_2008
10.1016/j.baae.2015.07.004	Ignasi_Bartomeus_Brassica_napus_Sweden_2013

10.1098/rspb.2020.0922	James_Reilly_Citrullus_lanatus_USA_2013, James_Reilly_Citrullus_lanatus_USA_2014, James_Reilly_Citrullus_lanatus_USA_2015, James_Reilly_Cucurbita_pepo_USA_2013, James_Reilly_Cucurbita_pepo_USA_2015, James_Reilly_Cucurbita_pepo_USA_2014, James_Reilly_Malus_pumila_USA_2013, James_Reilly_Malus_pumila_USA_2014, James_Reilly_Malus_pumila_USA_2015, James_Reilly_Prunus_avium_USA_2013, James_Reilly_Prunus_avium_USA_2014, James_Reilly_Prunus_cerasus_USA_2013, James_Reilly_Prunus_cerasus_USA_2014, James_Reilly_Prunus_cerasus_USA_2015, James_Reilly_Prunus_dulcis_USA_2013, James_Reilly_Prunus_dulcis_USA_2014, James_Reilly_Vaccinium_corymbosum_USA_2015, James_Reilly_Vaccinium_corymbosum_USA_2014, James_Reilly_Vaccinium_corymbosum_USA_2013
10.1111/1365-2664.12287	Jessica_D_Petersen_Cucurbita_pepo_USA_2011
10.1016/j.baae.2018.09.003	Jessica_Knapp_Cucurbita_pepo_UK_2016
10.1016/j.agee.2017.09.038	Juliana_Hipolito_Coffea_arabica_Brazil_2013, Juliana_Hipolito_Coffea_arabica_Brazil_2014

10.4257/oeco.2010.1401.09	Juliana_Hipolito_Mangifera_indica_Brazil_2005
10.3390/d12060259	Katrine_Hansen_Psidium_guajava_Thailand_2019, Katrine_Hansen_Psidium_guajava_Thailand_2020
10.1111/1365-2664.12977	Louis_Sutter_Brassica_napus_Switzerland_2014
10.1111/j.1461-0248.2010.01579.x	Luisa_G_Carvalho_Helianthus_annuus_South_Africa_2009
10.1111/j.1365-2664.2010.01829.x	Luisa_G_Carvalho_Mangifera_indica_South_Africa_2008
10.1111/j.1365-2664.2012.02217.x	Luisa_G_Carvalho_Mangifera_indica_South_Africa_2009
10.1007/s13592-018-0600-4	Marcos_Minarro_Malus_domestica_Spain_2015, Marcos_Minarro_Malus_domestica_Spain_2016
10.1017/CBO9780511754821	Margaret_Mayfield_Actinidia_deliciosa_New_Zealand_NA
10.1007/s10841-015-9788-z	Mark_Otieno_Cajanus_cajan_Kenya_2009
unpublished, 10.1016/j.biocon.2013.11.001	Michael_Garratt_Brassica_napus_UK_2012
unpublished, 10.1111/2041- 210X.13292	Michael_Garratt_Fragaria_ananassa_UK_2011
unpublished, 10.1371/journal.pone.0153889, 10.26786/1920- 7603(2014)8,10.1111/2041- 210X.13292	Michael_Garratt_Malus_domestica_UK_2011

unpublished, 10.1016/j.biocon.2013.11.001, 10.1111/2041-210X.13292	Michael_Garratt_Vicia_faba_UK_2011
10.1111/j.1365-2664.2005.01116.x, 10.1098/rspb.2007.1547	Natacha_Chacoff_Citrus_paradisi_Argentina_2000, Natacha_Chacoff_Citrus_paradisi_Argentina_2001, Natacha_Chacoff_Citrus_paradisi_Argentina_2002
https://hdl.handle.net/10214/21272	Leah_Blechschiidt_Malus_domestica_Canada_2018, Leah_Blechschiidt_Malus_domestica_Canada_2019
10.1111/j.1365-2664.2007.01418.x	Rachael_Winfree_Capsicum_annuum_USA_2004, Rachael_Winfree_Cucumis_melo_USA_2004, Rachael_Winfree_Solanum_lycopersicum_USA_2004, Rachael_Winfree_Solanum_lycopersicum_USA_2005
10.1111/j.1461-0248.2007.01110.x	Rachael_Winfree_Citrullus_lanatus_USA_2004, Rachael_Winfree_Citrullus_lanatus_USA_2005, Rachael_Winfree_Citrullus_lanatus_USA_2007, Rachael_Winfree_Citrullus_lanatus_USA_2008, Rachael_Winfree_Citrullus_lanatus_USA_2010, Rachael_Winfree_Citrullus_lanatus_USA_2011, Rachael_Winfree_Citrullus_lanatus_USA_2012
10.1111/1365-2664.12198	Rachael_Winfree_Vaccinium_corymbosum_USA_2010, Rachael_Winfree_Vaccinium_corymbosum_USA_2011
10.1111/ele.12126	Rachael_Winfree_Vaccinium_macrocarpon_USA_2009,

	Rachael_Winfree_Vaccinium_macrocarpon_USA_2010
10.1111/1365-2664.12377	Rachel_Mallinger_Malus_domestica_USA_2012, Rachel_Mallinger_Malus_domestica_USA_2013
10.1016/j.baae.2016.09.006	Rebecca_Steward_Fragaria_ananassa_Sweden_2014
10.1007/s00442-012-2271-6	Riccardo_Bommarco_Brassica_napus_Sweden_2005
10.1098/rspb.2011.0647	Riccardo_Bommarco_Trifolium_pratense_Sweden_2008, Riccardo_Bommarco_Trifolium_pratense_Sweden_2009, Riccardo_Bommarco_Trifolium_pratense_Sweden_2010
10.1007/s00442-015-3517-x	Sandra_Lindstrom_Brassica_napus_Sweden_2011, Sandra_Lindstrom_Brassica_napus_Sweden_2012
10.1016/j.agee.2016.04.020	Sarah_Cusser_Gossypium_hirsutum_USA_2014
10.1016/j.biocon.2006.05.025	Sarah_S_Greenleaf_Solanum_lycopersicum_USA_2001
10.1603/0022-0493-98.4.1193	Saul_A_Cunningham_Annona_squamosa atemoya_Australia_2001
10.1016/j.baae.2010.05.001	Saul_A_Cunningham_Brassica_napus_Australia_2006
10.1111/j.1600-0706.2009.17523.x	Shalene_Jha_Coffea_arabica_robusta_Mexico_2006
10.1016/j.baae.2012.03.007	Smitha_Krishnan_Coffea_canephora_India_2007, Smitha_Krishnan_Coffea_canephora_India_2008, Smitha_Krishnan_Coffea_canephora_India_2009
10.1073/pnas.0405147101,	Taylor_Ricketts_Coffea_arabica_Costa_Rica_2001,

10.1111/j.1523-1739.2004.00227.x	Taylor_Ricketts_Coffea_arabica_Costa_Rica_2002
10.1111/ele.13150	Thijs_Fijen_Allium_porrum_France_2016, Thijs_Fijen_Allium_porrum_Italy_2016
10.1007/s13593-016-0377-7, 10.1016/j.agee.2012.05.003, 10.1073/pnas.1210590110	Virginie_Boreux_Coffea_canephora_India_2008
10.1038/s41598-021-83341-7	Willis_Chan_Raine_Cucurbita_pepo_Canada_2017, Willis_Chan_Raine_Cucurbita_pepo_Canada_2018
10.1890/14-0910.1	Yael_Mandelik_Citrullus_lanatus_Israel_2009, Yael_Mandelik_Citrullus_lanatus_Israel_2010
10.1007/s13592-013-0242-5	Yael_Mandelik_Helianthus_annuus_Israel_2010
10.1186/s12898-017-0116-1	Yi_Zou_Brassica_napus_China_2015

LITERATURE CITED IN METADATA

Aizen, M. A., S. Aguiar, J. C. Biesmeijer, L. A. Garibaldi, D. W. Inouye, C. Jung, D. J. Martins,

R. Medel, C. L. Morales, H. Ngo, A. Pauw, R. J. Paxton, A. Sáez, & C. L. Seymour.

2019. Global agricultural productivity is threatened by increasing pollinator dependence without a parallel increase in crop diversification. *Global Change Biology*, 25(10), 3516–3527. <https://doi.org/10.1111/gcb.14736>

Bartomeus, I., and L. V. Dicks. 2019. The need for coordinated transdisciplinary research

infrastructures for pollinator conservation and crop pollination resilience. *Environmental Research Letters*, 14(4), 045017. <https://doi.org/10.1088/1748-9326/ab0cb5>

- Chamberlain, S., E. Szoecs, Z. Foster, Z. Arendsee, C. Boettiger, K. Ram, I. Bartomeus, J. Baumgartner, J. O'Donnell, J. Oksanen, B. Greshake Tzovaras, P. Marchand, V. Tran, M. Salmon, G. Li, and Grenié. 2020. *taxize: Taxonomic information from around the web*. R package version 0.9.98 <https://github.com/ropensci/taxize>.
- Dainese, M., E. A. Martin, M. A. Aizen, M. Albrecht, I. Bartomeus, R. Bommarco, L. G. Carvalheiro, R. Chaplin-Kramer, V. Gagic, L. A. Garibaldi, J. Ghazoul, H. Grab, M. Jonsson, D. S. Karp, C. M. Kennedy, D. Kleijn, C. Kremen, D. A. Landis, D. K. Letourneau, ... I. Steffan-Dewenter. 2019. A global synthesis reveals biodiversity-mediated benefits for crop production. *Science Advances*, 5(10), eaax0121. <https://doi.org/10.1126/sciadv.aax0121>
- Díaz, S., U. Pascual, M. Stenseke, B. Martín-López, R. T. Watson, Z. Molnár, R. Hill, K. M. A. Chan, I. A. Baste, K. A. Brauman, S. Polasky, A. Church, M. Lonsdale., A. Larigauderie, P. W. Leadley, A. P. E. van Oudenhoven, F. van der Plaats, M. Schröter, S. Lavorel, ... Y. Shirayama. 2018. Assessing nature's contributions to people. *Science*, 359(6373), 270–272. <https://doi.org/10.1126/science.aap8826>
- FAOSTAT data. 2018. Data available at <http://www.fao.org/faostat/en/#data/QC> Last accessed in November 2020.
- Garibaldi, L. A., I. Steffan-Dewenter, C. Kremen, J. M. Morales, R. Bommarco, S. A. Cunningham, L. G. Carvalheiro, N. P. Chacoff, J. H. Dudenhöffer, S. S. Greenleaf, A. Holzschuh, R. Isaacs, K. Krewenka, Y. Mandelik, M. M. Mayfield, L. A. Morandin, S. G. Potts, T. H. Ricketts, H. Szentgyörgyi, ... A.-M. Klein. 2011. Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters*, 14(10), 1062–1072. <https://doi.org/10.1111/j.1461-0248.2011.01669.x>

- Garibaldi, L. A., I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A. Cunningham, C. Kremen, L. G. Carvalheiro, L. D. Harder, O. Afik, I. Bartomeus, F. Benjamin, V. Boreux, D. Cariveau, N. P. Chacoff, J. H. Dudenhoffer, B. M. Freitas, J. Ghazoul, S. Greenleaf, ... A.-M. Klein. 2013. Wild Pollinators Enhance Fruit Set of Crops Regardless of Honey Bee Abundance. *Science*, 339(6127), 1608–1611.
<https://doi.org/10.1126/science.1230200>
- Garibaldi, L. A., I. Bartomeus, R. Bommarco, A.-M. Klein, S. A. Cunningham, M. A. Aizen, V. Boreux, M. P. D. Garratt, L. G. Carvalheiro, C. Kremen, C. L. Morales, C. Schüepp, N. P. Chacoff, B. M. Freitas, V. Gagic, A. Holzschuh, B. K. Klatt, K. M. Krewenka, S. Krishnan, ... M. Woyciechowski. 2015. EDITOR’S CHOICE: REVIEW: Trait matching of flower visitors and crops predicts fruit set better than trait diversity. *Journal of Applied Ecology*, 52(6), 1436–1444. <https://doi.org/10.1111/1365-2664.12530>
- Garibaldi, L. A., L. G. Carvalheiro, B. E. Vaissiere, B. Gemmill-Herren, J. Hipolito, B. M. Freitas, H. T. Ngo, N. Azzu, A. Saez, J. Astrom, J. An, B. Blochtein, D. Buchori, F. J. C. Garcia, F. Oliveira da Silva, K. Devkota, M. d. F. Ribeiro, L. Freitas, M. C. Gaglianone, ... H. Zhang. 2016. Mutually beneficial pollinator diversity and crop yield outcomes in small and large farms. *Science*, 351(6271), 388–391.
<https://doi.org/10.1126/science.aac7287>
- Garibaldi, L. A., A. Sáez, M. A. Aizen, T. Fijen, and I. Bartomeus. 2020. Crop pollination management needs flower-visitor monitoring and target values. *Journal of Applied Ecology*, 57(4), 664–670. <https://doi.org/10.1111/1365-2664.13574>
- Hampton, S. E., S. S. Anderson, S. C. Bagby, C. Gries, X. Han, E. M. Hart, M. B. Jones, W. C. Lenhardt, A. MacDonald, W. K. Michener, J. Mudge, A. Pourmokhtarian, M. P.

- Schildhauer, K. H. Woo, and N. Zimmerman. 2015. The Tao of open science for ecology. *Ecosphere*, 6(7), art120. <https://doi.org/10.1890/es14-00402.1>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, IPBES. 2016. Assessment Report on Pollinators, Pollination and Food Production. S.G. Potts, V. L. Imperatriz-Fonseca, and H. T. Ngo (eds). Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany. 552 pages. Zenodo. <https://doi.org/10.5281/ZENODO.3402856>
- Kleijn, D., R. Winfree, I. Bartomeus, L. G. Carvalheiro, M. Henry, R. Isaacs, A.-M. Klein, C. Kremen, L. K. M'Gonigle, R. Rader, T. H. Ricketts, N. M. Williams, N. Lee Adamson, J. S. Ascher, A. Báldi, P. Batáry, F. Benjamin, J. C. Biesmeijer, E. J. Blitzer, ... S. G. Potts. 2015. Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications*, 6, 7414. <https://doi.org/10.1038/ncomms8414>
- Klein, A.-M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, and T. Tscharntke. 2006. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608), 303–313. <https://doi.org/10.1098/rspb.2006.3721>
- Klein Goldewijk, K., A. Beusen, J. Doelman, and E. Stehfest. 2017. Anthropogenic land use estimates for the Holocene – HYDE 3.2. *Earth System Science Data*, 9(2), 927–953. <https://doi.org/10.5194/essd-9-927-2017>
- Millenium Ecosystem Assessment (MEA). 2005. *Ecosystems and human well being: synthesis*. Island Press, Washington, D.C., USA. <https://www.millenniumassessment.org/documents/document.356.aspx.pdf>

- Page, M. L., C. C. Nicholson, R. M. Brennan, A. T. Britzman, J. Greer, J. Hemberger, H. Kahl, U. Müller, Y. Peng, N. M. Rosenberger, C. Stuligross, L. Wang, L. H. Yang, & N. M. Williams. 2021. A meta-analysis of single visit pollination effectiveness. Cold Spring Harbor Laboratory. <https://doi.org/10.1101/2021.03.12.432378>
- Portman, Z. M., B. Bruninga-Socular, and D. P. Cariveau. 2020. The State of Bee Monitoring in the United States: A Call to Refocus Away From Bowl Traps and Towards More Effective Methods. *Annals of the Entomological Society of America*, 113(5), 337–342. <https://doi.org/10.1093/aesa/saaa010>
- Rader, R., I. Bartomeus, L. A. Garibaldi, M. P. D. Garratt, B. G. Howlett, R. Winfree, S. A. Cunningham, M. M. Mayfield, A. D. Arthur, G. K. S. Andersson, R. Bommarco, C. Brittain, L. G. Carvalheiro, N. P. Chacoff, M. H. Entling, B. Foully, B. M. Freitas, B. Gemmill-Herren, J. Ghazoul, ... M. Woyciechowski. 2015. Non-bee insects are important contributors to global crop pollination. *Proceedings of the National Academy of Sciences*, 113(1), 146–151. <https://doi.org/10.1073/pnas.1517092112>
- R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>
- Reilly, J. R., D. R. Artz, D. Biddinger, K. Bobiwash, N. K. Boyle, C. Brittain, J. Brokaw, J. W. Campbell, J. Daniels, E. Elle, J. D. Ellis, S. J. Fleischer, J. Gibbs, R. L. Gillespie, K. B. Gundersen, L. Gut, G. Hoffman, N. Joshi, O. Lundin, ... R. Winfree. 2020. Crop production in the USA is frequently limited by a lack of pollinators. *Proceedings of the Royal Society B: Biological Sciences*, 287(1931), 20200922. <https://doi.org/10.1098/rspb.2020.0922>

- Tscharntke, T., A.-M. Klein, A. Kruess, I. Steffan-Dewenter, and C. Thies. 2005. Landscape perspectives on agricultural intensification and biodiversity - ecosystem service management. *Ecology Letters*, 8(8), 857–874. <https://doi.org/10.1111/j.1461-0248.2005.00782.x>
- Vázquez, D. P., W. F. Morris, & P. Jordano. 2005. Interaction frequency as a surrogate for the total effect of animal mutualists on plants. *Ecology Letters*, 8(10), 1088–1094. <https://doi.org/10.1111/j.1461-0248.2005.00810.x>
- Venter, O., E. W. Sanderson, A. Magrach, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham, W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. M. Watson. 2016. Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications*, 7, 12558. <https://doi.org/10.1038/ncomms12558>
- White, E., E. Baldrige, Z. Brym, K. Locey, D. McGlinn, and S. Supp. 2013. Nine simple ways to make it easier to (re)use your data. *Ideas in Ecology and Evolution*, 6(2). <https://doi.org/10.4033/iee.2013.6b.6.f>
- Wickham, H. 2011. Testthat: Get started with testing. *The R Journal*, 3, 5–10. Retrieved from: http://journal.r-project.org/archive/2011-1/RJournal_2011-1_Wickham.pdf
- Winfree, R., J. R. Reilly, I. Bartomeus, D. P. Cariveau, N. M. Williams, and J. Gibbs. 2018. Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science*, 359(6377), 791–793. <https://doi.org/10.1126/science.aao2117>